

FOURIER–MUKAI FUNCTORS IN THE SUPPORTED CASE

ALBERTO CANONACO AND PAOLO STELLARI

ABSTRACT. We prove that exact functors between the categories of perfect complexes supported on projective schemes are of Fourier–Mukai type if the functor satisfies a condition weaker than being fully faithful. We also get generalizations of the results in the literature in the case without support conditions. Some applications are discussed and, along the way, we prove that the category of perfect supported complexes has a strongly unique enhancement.

1. INTRODUCTION

One of the most intriguing open questions in the theory of derived categories is whether all exact functors between the categories of perfect complexes (or between the bounded derived categories of coherent sheaves) on projective schemes are of Fourier–Mukai type. It might be worth recalling that, if X_1 and X_2 are projective schemes, an exact functor $F: \mathbf{Perf}(X_1) \rightarrow \mathbf{Perf}(X_2)$ between the corresponding categories of perfect complexes is a *Fourier–Mukai functor* (or of *Fourier–Mukai type*) if there exist $\mathcal{E} \in D^b(X_1 \times X_2)$ and an isomorphism of exact functors $F \cong \Phi_{\mathcal{E}}$. Here $\Phi_{\mathcal{E}}: \mathbf{Perf}(X_1) \rightarrow \mathbf{Perf}(X_2)$ is the exact functor defined by

$$\Phi_{\mathcal{E}} := \mathbf{R}(p_2)_*(\mathcal{E} \otimes^{\mathbf{L}} p_1^*(-)),$$

where $p_i: X_1 \times X_2 \rightarrow X_i$ is the natural projection. The complex \mathcal{E} is called a *kernel* of F .

While, in general, the kernel is certainly not unique (up to isomorphism) due to [10], the question about the existence of such kernels is widely open. Indeed, despite the fact that a conjecture in [4] would suggest a positive answer to it, for the time being, only partial results in this direction are available. Let us recall some of them. In [30] (together with [5]) the case of exact fully faithful functors between the bounded derived categories of coherent sheaves on smooth projective varieties is completely solved by Orlov. Various generalizations to quotient stacks and twisted categories were given in [19] by Kawamata and in [11] respectively. In particular, the main result of [11] shows that all exact functors $F: D^b(X_1) \rightarrow D^b(X_2)$ such that

$$(1.1) \quad \mathrm{Hom}_{D^b(X_2)}(F(\mathcal{A}), F(\mathcal{B})[k]) = 0,$$

for any $\mathcal{A}, \mathcal{B} \in \mathbf{Coh}(X_1)$ and any integer $k < 0$, are Fourier–Mukai functors and their kernels are unique, up to isomorphism.

The inspiration for our results in this paper comes from the new approach to the representability problem in [28], where the authors show that all exact fully faithful functors $F: \mathbf{Perf}(X_1) \rightarrow$

1991 *Mathematics Subject Classification.* 14F05, 18E10, 18E30.

Key words and phrases. Derived categories, Supported sheaves, Fourier–Mukai functors.

A. C. is partially supported by the national research project “Moduli, strutture geometriche e loro applicazioni” (PRIN 2009). P. S. is partially supported by the grants FIRB 2012 “Moduli Spaces and Their Applications” and the national research project “Geometria delle Varietà Proiettive” (PRIN 2010-11).

$\mathbf{Perf}(X_2)$ between the categories of perfect complexes on the projective schemes X_1 and X_2 are of Fourier–Mukai type. To show this, Lunts and Orlov prove that such fully faithful functors admit dg lifts. At that point, they can invoke the representability result in [32]. Indeed, Toën proved that, in the dg setting, all morphisms in the localization of the category of dg categories by quasi-equivalences are of Fourier–Mukai type (in an appropriate dg sense). Notice that the strategy in [28] allows the authors to improve the results in [1].

To make clear the categorical setting we are going to work with, let X_1 be a quasi-projective scheme containing a projective subscheme Z_1 such that the structure sheaf \mathcal{O}_{iZ_1} of the i -th infinitesimal neighbourhood of Z_1 in X_1 is in $\mathbf{Perf}(X_1)$, for every $i > 0$. This last condition is verified for instance when either $Z_1 = X_1$ or X_1 is smooth. Moreover let X_2 be a separated scheme of finite type over the base field \mathbb{k} with a closed subscheme Z_2 . One can then consider the categories $\mathbf{Perf}_{Z_i}(X_i)$ of perfect complexes on X_i with cohomology sheaves supported on Z_i . The definition of Fourier–Mukai functor makes perfect sense also in this context (see Definition 2.5).

A rewriting of (1.1) in the supported setting which weakens the fully-faithfulness condition in [28, 30] requires a bit of care. Indeed, assuming X_1, X_2, Z_1 and Z_2 to be as above, one can consider exact functors $F: \mathbf{Perf}_{Z_1}(X_1) \rightarrow \mathbf{Perf}_{Z_2}(X_2)$ such that

- (1) $\mathrm{Hom}(F(\mathcal{A}), F(\mathcal{B})[k]) = 0$, for any $\mathcal{A}, \mathcal{B} \in \mathbf{Coh}_{Z_1}(X_1) \cap \mathbf{Perf}_{Z_1}(X_1)$ and any integer $k < 0$;
- (*) (2) For all $\mathcal{A} \in \mathbf{Perf}_{Z_1}(X_1)$ with trivial cohomologies in positive degrees, there is $N \in \mathbb{Z}$ such that

$$\mathrm{Hom}(F(\mathcal{A}), F(\mathcal{O}_{|Z_1}(jH_1))) = 0,$$

for any $i < N$ and any $j \ll i$, where H_1 is an ample (Cartier) divisor on X_1 .

At first sight this condition may look a bit involved, but if $Z_1 = X_1$ is smooth with $\dim(X_1) > 0$, then part (2) of (*) is redundant and thus (*) turns out to be equivalent to (1.1) (see Proposition 3.13). In general full functors always satisfy (*), if we assume further that the maximal 0-dimensional torsion subsheaf $T_0(\mathcal{O}_{Z_1})$ of \mathcal{O}_{Z_1} is trivial. Actually, due to [7], a non-trivial full functor is automatically faithful if Z_1 is connected. We will discuss in Section 3.4 the existence of non-full functors with property (*).

We are now ready to state our first main result.

Theorem 1.1. *Let X_1 be a quasi-projective scheme containing a projective subscheme Z_1 such that $\mathcal{O}_{iZ_1} \in \mathbf{Perf}(X_1)$, for all $i > 0$, and let X_2 be a separated scheme of finite type over the base field \mathbb{k} with a closed subscheme Z_2 . Let*

$$F: \mathbf{Perf}_{Z_1}(X_1) \longrightarrow \mathbf{Perf}_{Z_2}(X_2)$$

be an exact functor.

If F satisfies (*), then there exist $\mathcal{E} \in D_{Z_1 \times Z_2}^b(\mathbf{Qcoh}(X_1 \times X_2))$ and an isomorphism of functors $F \cong \Phi_{\mathcal{E}}^s$. Moreover, if X_i is smooth quasi-projective, for $i = 1, 2$, and \mathbb{k} is perfect, then \mathcal{E} is unique up to isomorphism.

This result summarizes the content of Proposition 4.10, Lemma 5.3 and Proposition 5.6. The proof is contained in Sections 4 and 5 and it uses the approach via dg categories proposed in [28].

Clearly, assuming $X_i = Z_i$ for $i = 1, 2$, our result extends the one in [28] about singular projective schemes (see Corollary 4.11). Notice that the symbol $\Phi_{\mathcal{E}}^s$ stands for the ‘supported’ Fourier–Mukai functor with kernel \mathcal{E} defined precisely in (2.2).

Our second main result concerns the uniqueness of the enhancement for the category of perfect supported complexes mentioned above and it is proved in Section 4.2.

Theorem 1.2. *Let X be a quasi-projective scheme containing a projective subscheme Z such that $\mathcal{O}_{iZ} \in \mathbf{Perf}(X)$, for all $i > 0$, and $T_0(\mathcal{O}_Z) = 0$. Then $\mathbf{Perf}_Z(X)$ has a strongly unique enhancement.*

The notion of enhancement and its strong uniqueness is discussed in Section 4.1. For the moment we can roughly think of an enhancement of $\mathbf{Perf}_Z(X)$ as a (pretriangulated) dg category whose homotopy category is equivalent to $\mathbf{Perf}_Z(X)$. The enhancement is strongly unique if two such are (quasi-)equivalent at the dg category level and such an equivalence satisfies some additional condition. It is worth noticing that the particular case $X = Z$ is one of the main results in [28] (see Corollary 4.9).

Motivations. Due to the technical nature of Theorems 1.1 and 1.2, some geometric motivations are certainly in order here. From our point of view the reason for studying exact functors between categories with support conditions is twofold. On one side the conjecture in [4] concerning the fact that all ‘geometric’ functors are of Fourier–Mukai type appears extremely difficult to prove in complete generality. Thus it makes sense to test its validity by weakening the assumptions on the geometric nature of the triangulated categories involved and on the exact functors between them. In this sense, this paper is in the same spirit as [11] and [28].

On the other hand, one would like to study easy-to-handle *d-Calabi–Yau categories*, i.e. triangulated categories whose Serre functor is isomorphic to the shift by the positive integer d . Challenging examples are certainly provided by the derived categories of smooth projective Calabi–Yau threefolds. Indeed, the homological version of the Mirror Symmetry conjecture [24] for those threefolds involves these categories with implications for the manifolds parametrizing stability conditions [6] into which (up to the quotient by the group of autoequivalences) the Kähler moduli spaces embed. One big open problem in this direction is the lack of examples of stability conditions for Calabi–Yau threefolds.

The group of autoequivalences of the derived category, besides being an interesting algebraic object in itself, acts on the stability manifold. Already for Calabi–Yau manifolds of dimension 2 (i.e. K3 surfaces), this group is very complicated and one of the main motivations of [30] is to stimulate its study. As for stability conditions, in higher dimension the situation becomes much more involved.

Therefore, following suggestions from the physics literature, one may start from the non-compact or the so called ‘open’ Calabi–Yau manifolds. Let us be more precise discussing some explicit examples where the ambient space X_1 is smooth and Theorem 1.1 (or a variant of it) applies.

Following [13] and [23], one can consider the triangulated category \mathbf{T}_S classically generated by a d -spherical object S (here d is a positive integer) in an idempotent complete triangulated category \mathbf{T} . An object S is d -spherical if the graded algebra $\mathrm{Ext}^*(S, S)$ is isomorphic to the cohomology of

a d -sphere. We will study this example in Section 4.4 when $d = 1$ as in this case \mathbf{T}_S is nothing but $D_{\mathbb{P}^1}^b(C)$, where C is a smooth curve and $p \in C$ is a \mathbb{k} -rational point. Thus we obtain the following result, which is a particular case of Proposition 4.15.

Proposition 1.3. *Every exact autoequivalence of \mathbf{T}_S is of Fourier–Mukai type if S is a 1-spherical object.*

This completes the picture in [13] which provides a description of the subgroup of Fourier–Mukai autoequivalences. We should remark here that the result above is not a direct consequence of Theorem 1.1 as the maximal 0-dimensional torsion subsheaf of \mathcal{O}_p is obviously not trivial and part (2) of (*) does not hold true.

Interesting examples of 2-Calabi–Yau categories are provided by the local resolutions of A_n -singularities on surfaces which were studied in [17, 16]. More precisely, one considers $Y = \text{Spec}(\mathbb{C}[[x, y, z]]/(x^2 + y^2 + z^{n+1}))$ (the A_n -singularity), the minimal resolution $f: X \rightarrow Y$ and $Z := f^{-1}(p)$, where p is the closed point in Y . Notice that, in this case, $T_0(\mathcal{O}_Z) = 0$. The category one wants to consider is then $D_Z^b(X) = \mathbf{Perf}_Z(X)$ and using Theorem 1.1 we can reprove in a direct way the following result already contained in [16].

Corollary 1.4. *Every exact autoequivalence of $D_Z^b(X)$ is of Fourier–Mukai type.*

Finally, to get examples of 3-Calabi–Yau categories one can take the total space $\text{tot}(\omega_{\mathbb{P}^2})$ of the canonical bundle of \mathbb{P}^2 . In this case, if Z denotes the zero section of the projection $\text{tot}(\omega_{\mathbb{P}^2}) \rightarrow \mathbb{P}^2$, the derived category $\mathbf{Perf}_Z(\text{tot}(\omega_{\mathbb{P}^2})) = D_Z^b(\text{tot}(\omega_{\mathbb{P}^2}))$ is a 3-Calabi–Yau category and may be seen as an interesting example to test predictions about Mirror Symmetry and the topology of the space of stability conditions according to Bridgeland’s definition (see [3] for results in this direction). Here again $T_0(\mathcal{O}_Z) = 0$ and so Theorem 1.1 yields the following.

Corollary 1.5. *Every exact autoequivalence of $D_Z^b(\text{tot}(\omega_{\mathbb{P}^2}))$ is of Fourier–Mukai type.*

As an application of Proposition 4.15 and Theorem 1.2, the triangulated categories in the three examples above have strongly unique enhancements.

The plan of the paper. In Section 2 we provide the necessary preliminary material concerning derived categories of supported sheaves, and we introduce the notion of an almost ample set. Then we prove a criterion (generalizing others present in the literature) for extending a morphism defined on a suitable subset of the source category between exact functors satisfying a condition related to (*). This is done in Section 3 using the notion of convolution. In Section 4 we deal with the existence of Fourier–Mukai kernels and the strong uniqueness of enhancements. In particular, we need to generalize and to modify the argument in [28] to make it work in our setting. In the same section we also discuss the case of 1-spherical objects. Section 5 deals with various questions about boundedness and uniqueness of Fourier–Mukai kernels.

Notation. In this paper, \mathbb{k} is a field. All schemes are assumed to be at least of finite type and separated over \mathbb{k} . All additive (in particular, triangulated) categories and all additive (in particular, exact) functors will be assumed to be \mathbb{k} -linear. If \mathbf{A} is an abelian (or more generally an exact)

category, $D(\mathbf{A})$ denotes the derived category of \mathbf{A} and $D^b(\mathbf{A})$ its full subcategory of complexes with bounded cohomology. For an object

$$A := \{ \cdots \rightarrow A^j \xrightarrow{d^j} A^{j+1} \xrightarrow{d^{j+1}} \cdots \xrightarrow{d^{i-1}} A^i \xrightarrow{d^i} A^{i+1} \rightarrow \cdots \}$$

in $D(\mathbf{A})$, we can consider the *gentle truncations* $\tau_{\leq i}A$, $\tau_{\geq i}A$, defined as

$$\begin{aligned} \tau_{\leq i}A &:= \{ \cdots \rightarrow A^j \xrightarrow{d^j} A^{j+1} \xrightarrow{d^{j+1}} \cdots \xrightarrow{d^{i-1}} \ker d^i \rightarrow 0 \rightarrow \cdots \} \\ \tau_{\geq i}A &:= \{ \cdots \rightarrow 0 \rightarrow \operatorname{coker} d^{i-1} \rightarrow A^{i+1} \xrightarrow{d^{i+1}} \cdots \rightarrow A^j \xrightarrow{d^j} \cdots \}. \end{aligned}$$

Unless clearly stated, all functors are derived even if, for simplicity, we use the same symbol for a functor and its derived version. Natural transformations (in particular, isomorphisms) between exact functors are always assumed to be compatible with shifts.

2. PRELIMINARIES

The first part of this section provides a quick introduction to some basic and well-known facts concerning the derived categories of supported sheaves. Then we define and discuss the notion of an almost ample set.

2.1. Categories with support conditions. Let X be a scheme and let Z be a closed subscheme of X . We denote by $D_Z(\mathbf{Qcoh}(X))$ the derived category of unbounded complexes of quasi-coherent sheaves on X with cohomologies supported on Z . We will be particularly interested in the triangulated categories

$$(2.1) \quad \begin{aligned} D_Z^b(\mathbf{Qcoh}(X)) &:= D_Z(\mathbf{Qcoh}(X)) \cap D^b(\mathbf{Qcoh}(X)) \\ D_Z^b(X) &:= D_Z(\mathbf{Qcoh}(X)) \cap D^b(X), \end{aligned}$$

where $D^b(X) := D_{\mathbf{Coh}}^b(\mathbf{Qcoh}(X))$ is the full subcategory of $D^b(\mathbf{Qcoh}(X))$ consisting of complexes with coherent cohomologies. Denote by $\mathbf{Perf}(X) \subset D(\mathbf{Qcoh}(X))$ the full subcategory of perfect complexes on X . Notice that $\mathbf{Perf}(X) \subseteq D^b(X)$ and, if X is quasi-projective, equality holds if and only if X is regular. In the supported case we set

$$\mathbf{Perf}_Z(X) := D_Z(\mathbf{Qcoh}(X)) \cap \mathbf{Perf}(X).$$

Thus, if X is smooth, $\mathbf{Perf}_Z(X) = D_Z^b(X)$.

Proposition 2.1. ([31], **Theorems 5.3(i) and 6.8.**) *The category $D_Z(\mathbf{Qcoh}(X))$ is compactly generated and the subcategory of compact objects $D_Z(\mathbf{Qcoh}(X))^c$ coincides with $\mathbf{Perf}_Z(X)$.*

Recall that an object A in a triangulated category \mathbf{T} is *compact* if, given $\{X_i\}_{i \in I} \subset \mathbf{T}$ such that I is a set and $\bigoplus_i X_i$ exists in \mathbf{T} , the canonical map

$$\bigoplus_i \operatorname{Hom}(A, X_i) \longrightarrow \operatorname{Hom}(A, \bigoplus_i X_i)$$

is an isomorphism. Moreover, \mathbf{T} is *compactly generated* if there is a set S of objects in the full subcategory \mathbf{T}^c of compact objects of \mathbf{T} such that, given $E \in \mathbf{T}$ with $\operatorname{Hom}(A, E[i]) = 0$ for all $A \in S$ and all $i \in \mathbb{Z}$, then $E \cong 0$. For more details, the reader can consult [31, Sect. 3.1].

The category $D_Z(\mathbf{Qcoh}(X))$ is a full subcategory of $D(\mathbf{Qcoh}(X))$ and let

$$\iota: D_Z(\mathbf{Qcoh}(X)) \hookrightarrow D(\mathbf{Qcoh}(X))$$

be the inclusion. We use the same symbol to denote the inclusion functor for the other categories in (2.1). According to [27, Sect. 3], the functor ι has a right adjoint

$$\iota^!: D(\mathbf{Qcoh}(X)) \rightarrow D_Z(\mathbf{Qcoh}(X)).$$

Notice that the existence of $\iota^!$ could be also deduced from [29], since ι clearly commutes with arbitrary direct sums. As ι is fully faithful, we have $\iota^! \circ \iota \cong \text{id}$.

Remark 2.2. Actually [27] deals only with modules over a commutative noetherian ring. On the other hand, as it is observed in the introduction of [27], most of its results globalize to sheaves over schemes (in particular those used in this section).

Lemma 2.3. *The functor $\iota^!$ sends bounded complexes to bounded complexes and commutes with direct sums.*

Proof. The fact that $\iota^!$ sends bounded complexes to bounded complexes follows from [27, Cor. 3.1.4]. The commutativity with direct sums is due to [27, Cor. 3.5.2]. \square

Now, let X_1 and X_2 be schemes containing, respectively, two closed subschemes Z_1 and Z_2 . We will denote by ι_i (respectively $\iota_{i,j}$) the inclusion morphisms relative to the pair (X_i, Z_i) (respectively $(X_i \times X_j, Z_i \times Z_j)$), for $i, j = 1, 2$.

Lemma 2.4. *Let $f: X_1 \rightarrow X_2$ be a morphism of schemes such that $f^{-1}(Z_2) = Z_1$. Then there are isomorphisms of exact functors*

$$\begin{aligned} f_* \circ \iota_1 \circ \iota_1^! &\cong \iota_2 \circ \iota_2^! \circ f_*: D(\mathbf{Qcoh}(X_1)) \rightarrow D(\mathbf{Qcoh}(X_2)) \\ f^* \circ \iota_2 \circ \iota_2^! &\cong \iota_1 \circ \iota_1^! \circ f^*: D(\mathbf{Qcoh}(X_2)) \rightarrow D(\mathbf{Qcoh}(X_1)). \end{aligned}$$

Moreover, given a pair (X, Z) , for every $\mathcal{F} \in D(\mathbf{Qcoh}(X))$ there is an isomorphism of exact functors

$$\iota \circ \iota^!(\mathcal{F} \otimes -) \cong \mathcal{F} \otimes \iota \circ \iota^!(-): D(\mathbf{Qcoh}(X)) \rightarrow D(\mathbf{Qcoh}(X)).$$

Proof. See Cor. 3.4.3, 3.4.4 and 3.3.1 of [27]. \square

Definition 2.5. An exact functor

$$F: D_{Z_1}(\mathbf{Qcoh}(X_1)) \rightarrow D_{Z_2}(\mathbf{Qcoh}(X_2))$$

is a *Fourier–Mukai functor* if there exist $\mathcal{E} \in D_{Z_1 \times Z_2}(\mathbf{Qcoh}(X_1 \times X_2))$ and an isomorphism of exact functors

$$(2.2) \quad F \cong \Phi_{\mathcal{E}}^s := \iota_2^! \circ (p_2)_* (\iota_{1,2}(\mathcal{E}) \otimes p_1^* \circ \iota_1(-))$$

where $p_i: X_1 \times X_2 \rightarrow X_i$ is the projection.

Analogous definitions can be given for functors defined between bounded derived categories of quasi-coherent, coherent or perfect complexes. The object \mathcal{E} is called *Fourier–Mukai kernel*. We will use the standard notation $\Phi_{\mathcal{E}}$ when $Z_i = X_i$ or to denote Fourier–Mukai functors between $D(\mathbf{Qcoh}(X_1))$ and $D(\mathbf{Qcoh}(X_2))$.

Observe that, as $(p_2)_*(\iota_{1,2}(\mathcal{E}) \otimes p_1^* \circ \iota_1(-))$ is supported on Z_2 , $\iota_2 \circ \Phi_{\mathcal{E}}^s \cong \Phi_{\iota_{1,2}(\mathcal{E})} \circ \iota_1$. This clearly implies that $\Phi_{\mathcal{E}}^s \cong \iota_2^! \circ \Phi_{\iota_{1,2}(\mathcal{E})} \circ \iota_1$, and more generally we can prove the following result.

Lemma 2.6. *Under the above assumptions, for every $\tilde{\mathcal{E}} \in D(\mathbf{Qcoh}(X_1 \times X_2))$ such that $\mathcal{E} \cong \iota_{1,2}^!(\tilde{\mathcal{E}})$ there exists an isomorphism of exact functors*

$$\Phi_{\mathcal{E}}^s \cong \iota_2^! \circ \Phi_{\tilde{\mathcal{E}}} \circ \iota_1 : D_{Z_1}(\mathbf{Qcoh}(X_1)) \rightarrow D_{Z_2}(\mathbf{Qcoh}(X_2)).$$

Proof. Denoting by $\bar{\iota}_1$ and $\bar{\iota}_2$ the inclusion morphisms relative to the pairs $(X_1 \times X_2, Z_1 \times X_2)$ and $(X_1 \times X_2, X_1 \times Z_2)$ respectively, it is easy to see that $\iota_{1,2} \circ \iota_{1,2}^! \cong \bar{\iota}_2 \circ \bar{\iota}_2^! \circ \bar{\iota}_1 \circ \bar{\iota}_1^!$. Therefore, using Lemma 2.4 repeatedly, we obtain

$$\begin{aligned} \Phi_{\mathcal{E}}^s &\cong \iota_2^! \circ (p_2)_*(\iota_{1,2} \circ \iota_{1,2}^!(\tilde{\mathcal{E}}) \otimes p_1^* \circ \iota_1(-)) \cong \iota_2^! \circ (p_2)_*(\bar{\iota}_2 \circ \bar{\iota}_2^! \circ \bar{\iota}_1 \circ \bar{\iota}_1^!(\tilde{\mathcal{E}}) \otimes p_1^* \circ \iota_1(-)) \\ &\cong \iota_2^! \circ (p_2)_* \circ \bar{\iota}_2 \circ \bar{\iota}_2^!(\bar{\iota}_1 \circ \bar{\iota}_1^!(\tilde{\mathcal{E}}) \otimes p_1^* \circ \iota_1(-)) \cong \iota_2^! \circ \iota_2 \circ \iota_2^! \circ (p_2)_*(\bar{\iota}_1 \circ \bar{\iota}_1^!(\tilde{\mathcal{E}}) \otimes p_1^* \circ \iota_1(-)), \end{aligned}$$

whence

$$\begin{aligned} \Phi_{\mathcal{E}}^s \circ \iota_1^! &\cong \iota_2^! \circ (p_2)_*(\bar{\iota}_1 \circ \bar{\iota}_1^!(\tilde{\mathcal{E}}) \otimes p_1^* \circ \iota_1 \circ \iota_1^!(-)) \cong \iota_2^! \circ (p_2)_*(\bar{\iota}_1 \circ \bar{\iota}_1^!(\tilde{\mathcal{E}}) \otimes \bar{\iota}_1 \circ \bar{\iota}_1^! \circ p_1^*(-)) \\ &\cong \iota_2^! \circ (p_2)_*(\tilde{\mathcal{E}} \otimes \bar{\iota}_1 \circ \bar{\iota}_1^! \circ p_1^*(-)) \cong \iota_2^! \circ (p_2)_*(\tilde{\mathcal{E}} \otimes p_1^* \circ \iota_1 \circ \iota_1^!(-)) = \iota_2^! \circ \Phi_{\tilde{\mathcal{E}}} \circ \iota_1 \circ \iota_1^!. \end{aligned}$$

So we conclude that $\Phi_{\mathcal{E}}^s \cong \Phi_{\tilde{\mathcal{E}}} \circ \iota_1^! \circ \iota_1 \cong \iota_2^! \circ \Phi_{\tilde{\mathcal{E}}} \circ \iota_1 \circ \iota_1^! \cong \iota_2^! \circ \Phi_{\tilde{\mathcal{E}}} \circ \iota_1$. \square

Consider the abelian categories $\mathbf{Qcoh}_Z(X)$ and $\mathbf{Coh}_Z(X)$ consisting of quasi-coherent and, respectively, coherent sheaves supported on Z . The following will be implicitly used at many points of this paper.

Proposition 2.7. ([2], Lemma 3.3, Corollary 3.4.) *The natural functors $D(\mathbf{Qcoh}_Z(X)) \rightarrow D_Z(\mathbf{Qcoh}(X))$ and $D^b(\mathbf{Coh}_Z(X)) \rightarrow D_Z^b(X)$ are equivalences.*

Notice that the proof in [2] works in our generality as well. Denoting by $i: Z \hookrightarrow X$ the closed embedding, we will also need the following result.

Proposition 2.8. ([2], Lemma 3.6, Corollary 3.7.) (i) *The functor i_* is exact and the image of i_* generates $\mathbf{Coh}_Z(X)$ as an abelian category, i.e. the smallest abelian subcategory of $\mathbf{Coh}_Z(X)$ closed under extensions and containing the essential image of i_* is $\mathbf{Coh}_Z(X)$ itself. Similarly, the smallest abelian subcategory closed under extensions and arbitrary direct sums containing the image of i_* in $\mathbf{Qcoh}_Z(X)$ is $\mathbf{Qcoh}_Z(X)$ itself.*

(ii) *The image of $D^b(Z)$ under i_* classically generates $D_Z^b(X)$ and the image under i_* of $D(\mathbf{Qcoh}(Z))$ classically completely generates $D_Z(\mathbf{Qcoh}(X))$.*

Recall that, according to [31], a subcategory \mathbf{S} of a triangulated category \mathbf{T} classically generates \mathbf{T} if the smallest thick triangulated subcategory of \mathbf{T} containing \mathbf{S} is \mathbf{T} itself. On the other hand, \mathbf{S} classically completely generates \mathbf{T} if \mathbf{T} is the smallest thick subcategory which is closed under direct sums and contains \mathbf{S} .

2.2. Almost ample sets. The attempt of this section is to provide a generalization of the notion of weakly ample sequence in [16, Appendix A.2] which, in turn, is a generalization of the usual definition of ample sequence (see, for example, [30]).

Definition 2.9. Given an abelian category \mathbf{A} and a set I , a subset $\{P_i\}_{i \in I} \subseteq \mathbf{A}$ is an *almost ample set* if, for any $A \in \mathbf{A}$, there exists $i \in I$ such that:

- (1) there is a natural number k and an epimorphism $P_i^{\oplus k} \twoheadrightarrow A$;
- (2) $\mathrm{Hom}_{\mathbf{A}}(A, P_i) = 0$.

Remark 2.10. Every ample sequence and, more generally, every weakly ample sequence is an almost ample set (with $I = \mathbb{Z}$). It is also obvious by definition that every set containing an almost ample set (respectively a set of objects satisfying (1) in Definition 2.9) is almost ample (respectively it satisfies (1) in Definition 2.9), too.

To provide examples of almost ample sets which are suited for the supported setting we are working in, let X be a quasi-projective scheme and let Z be a projective subscheme of X . Assume further that $\mathcal{O}_{iZ} \in \mathbf{Perf}(X)$, for all $i > 0$. Take H an ample divisor on X and define the subset of $\mathbf{Coh}_Z(X)$

$$(2.3) \quad \mathbf{Amp}(Z, X, H) := \{\mathcal{O}_{|i|Z}(jH)\}_{(i,j) \in \mathbb{Z} \times \mathbb{Z}}.$$

When needed, we will think of $\mathbf{Amp}(Z, X, H)$ as the corresponding full subcategory of $\mathbf{Coh}_Z(X)$.

Example 2.11. There are two interesting geometric situations for which $\mathcal{O}_{iZ} \in \mathbf{Perf}(X)$, for all $i > 0$, and thus $\mathbf{Amp}(Z, X, H)$ is contained in $\mathbf{Perf}_Z(X)$. Namely one can take X to be a quasi-projective scheme containing a projective subscheme Z such that either $Z = X$ or X is smooth.

The following result will be essential for the rest of the paper. Using, for example, the notation of [15, Def. 1.1.4], we denote by $T_0(\mathcal{O}_Z)$ the maximal subsheaf of \mathcal{O}_Z whose support has dimension 0. For short, we call $T_0(\mathcal{O}_Z)$ the *maximal 0-dimensional torsion subsheaf* of \mathcal{O}_Z .

Proposition 2.12. *Assume that X , Z and H are as above. Then $\mathbf{Amp}(Z, X, H)$ satisfies (1) in Definition 2.9, and provides a set of (compact) generators of the Grothendieck category $\mathbf{Qcoh}_Z(X)$. Moreover, if $T_0(\mathcal{O}_Z) = 0$, then $\mathbf{Amp}(Z, X, H)$ is an almost ample set in $\mathbf{Coh}_Z(X)$.*

More precisely, for any $\mathcal{A} \in \mathbf{Coh}_Z(X)$, there is $N \in \mathbb{Z}$ such that any $\mathcal{O}_{|i|Z}(jH)$ with $i < N$ and $j \ll i$ satisfies (1) (and (2) if $T_0(\mathcal{O}_Z) = 0$) in Definition 2.9.

Proof. Notice that under the assumption $\mathcal{O}_{iZ} \in \mathbf{Perf}(X)$ for all $i > 0$, the objects $\mathcal{O}_{|i|Z}(jH)$ are compact as well for all $i, j \in \mathbb{Z}$.

Let \mathcal{E} be a sheaf in $\mathbf{Coh}_Z(X)$. To prove property (1), observe that by [31, Lemma 7.40] there is an integer $N \leq 0$ such that \mathcal{E} is a coherent $\mathcal{O}_{|i|Z}$ -module for all $i < N$. As $|i|Z$ is a projective scheme, it follows that for $i < N$ and $j \ll i$ there is an epimorphism $\mathcal{O}_{|i|Z}(jH)^{\oplus k} \twoheadrightarrow \mathcal{E}$ (for some $k \in \mathbb{N}$) in $\mathbf{Coh}(|i|Z)$, whence also in $\mathbf{Coh}_Z(X)$.

Assuming $T_0(\mathcal{O}_Z) = 0$ (which, indeed, implies $T_0(\mathcal{O}_{|i|Z}) = 0$), property (2) is easily verified taking $i < N$ and $j \ll i$. In fact, for $i < N$ and any $j \in \mathbb{Z}$, we have

$$\mathrm{Hom}_{\mathbf{Coh}_Z(X)}(\mathcal{E}, \mathcal{O}_{|i|Z}(jH)) \cong \mathrm{Hom}_{\mathbf{Coh}(|i|Z)}(\mathcal{E}, \mathcal{O}_{|i|Z}(jH)).$$

Under the assumption $T_0(\mathcal{O}_{|i|Z}) = 0$, the same argument as in the proof of [28, Prop. 9.2] yields that, for j sufficiently small, the latter vector space is trivial.

To prove that $\mathbf{Amp}(Z, X, H)$ is a set of generators for the category $\mathbf{Qcoh}_Z(X)$, it is enough to observe that, in view of (1) of Definition 2.9 and the fact that any quasi-coherent sheaf is the direct limit of its coherent subsheaves, for any $\mathcal{E} \in \mathbf{Qcoh}_Z(X)$ there is a surjection $\bigoplus_{j \in S} P_j \rightarrow \mathcal{E}$ where S is a set and $P_j \in \mathbf{Amp}(Z, X, H)$ for all $j \in S$ (see [18, Sect. 8.3]). \square

Example 2.13. If X is the resolution of an A_n -singularity and Z is the exceptional locus, a special case of almost ample set for $\mathbf{Coh}_Z(X)$ is provided by the *weak ample sequence* \mathbf{C} in [16, Appendix A], where $\mathbf{C} = \{\mathcal{O}_{|i|Z}(iH) \in \mathbf{Amp}(Z, X, H) : i \in \mathbb{Z}\}$.

3. EXTENDING NATURAL TRANSFORMATIONS

In this section we deal with the second key ingredient in our proof, namely a criterion to extend natural transformations (in particular, isomorphisms) between functors. We tried to put this result in a generality which goes beyond the scope of this paper but which may be useful in future works (see, for example, [9, Prop. 5.15]).

3.1. Convolutions. In this section we collect some well-known facts about convolutions which will be used in the paper. Most of the terminology is taken from [19, 30] (see also [11]).

A bounded complex in a triangulated category \mathbf{T} is a sequence of objects and morphisms in \mathbf{T}

$$(3.1) \quad A_m \xrightarrow{d_m} A_{m-1} \xrightarrow{d_{m-1}} \cdots \xrightarrow{d_1} A_0$$

such that $d_j \circ d_{j+1} = 0$ for $0 < j < m$. A *right convolution* of (3.1) is an object A together with a morphism $d_0: A_0 \rightarrow A$ such that there exists a diagram in \mathbf{T}

$$\begin{array}{ccccccccccc} A_m & \xrightarrow{d_m} & A_{m-1} & \xrightarrow{d_{m-1}} & \cdots & \xrightarrow{d_2} & A_1 & \xrightarrow{d_1} & A_0 & & \\ & \searrow \text{id} & \swarrow \circlearrowleft & \searrow \circlearrowleft & \swarrow \circlearrowleft & \searrow \circlearrowleft & \swarrow \circlearrowleft & \searrow \circlearrowleft & \swarrow \circlearrowleft & \searrow d_0 & \\ & & A_m & \xleftarrow{[1]} & C_{m-1} & \xleftarrow{[1]} & \cdots & \xleftarrow{[1]} & C_1 & \xleftarrow{[1]} & A \end{array}$$

where the triangles with a \circlearrowleft are commutative and the others are distinguished.

Let $d_0: A_0 \rightarrow A$ be a right convolution of (3.1). If \mathbf{T}' is another triangulated category and $\mathbf{G}: \mathbf{T} \rightarrow \mathbf{T}'$ is an exact functor, then $\mathbf{G}(d_0): \mathbf{G}(A_0) \rightarrow \mathbf{G}(A)$ is a right convolution of

$$\mathbf{G}(A_m) \xrightarrow{\mathbf{G}(d_m)} \mathbf{G}(A_{m-1}) \xrightarrow{\mathbf{G}(d_{m-1})} \cdots \xrightarrow{\mathbf{G}(d_1)} \mathbf{G}(A_0).$$

The following results will be used in the rest of this section. Notice that left convolutions in the notation of [19] correspond to right convolutions in the present paper (according to the notation in [11, 30]).

Lemma 3.1. ([19], Lemmas 2.1 and 2.4.) *Let (3.1) be a complex in \mathbf{T} satisfying*

$$(3.2) \quad \text{Hom}_{\mathbf{T}}(A_a, A_b[r]) = 0 \text{ for any } a > b \text{ and } r < 0.$$

Then (3.1) has a right convolution which is uniquely determined up to isomorphism (in general non-canonical).

Lemma 3.2. ([11], **Lemma 3.3.**) *Let*

$$\begin{array}{ccccccc}
 A_m & \xrightarrow{d_m} & A_{m-1} & \xrightarrow{d_{m-1}} & \cdots & \xrightarrow{d_2} & A_1 & \xrightarrow{d_1} & A_0 \\
 \downarrow f_m & & \downarrow f_{m-1} & & & & \downarrow f_1 & & \downarrow f_0 \\
 B_m & \xrightarrow{e_m} & B_{m-1} & \xrightarrow{e_{m-1}} & \cdots & \xrightarrow{e_2} & B_1 & \xrightarrow{e_1} & B_0
 \end{array}$$

be a morphism of complexes both satisfying (3.2) and such that

$$\mathrm{Hom}_{\mathbf{T}}(A_a, B_b[r]) = 0 \text{ for any } a > b \text{ and } r < 0.$$

Assume that the corresponding right convolutions are of the form $(d_0, 0): A_0 \rightarrow A \oplus \bar{A}$ and $(e_0, 0): B_0 \rightarrow B \oplus \bar{B}$ and that $\mathrm{Hom}_{\mathbf{T}}(A_p, B[r]) = 0$ for $r < 0$ and any p . Then there exists a unique morphism $f: A \rightarrow B$ such that $f \circ d_0 = e_0 \circ f_0$. If moreover each f_i is an isomorphism, then f is an isomorphism as well.

Let $\mathbf{T} := \mathrm{D}^b(\mathbf{A})$ for some abelian category \mathbf{A} and let E be a complex as in (3.1) and such that every A_i is an object of \mathbf{A} . Then a right convolution of E (which is unique up to isomorphism by Lemma 3.1) is the natural morphism $A_0 \rightarrow E^\bullet$, where E^\bullet is the object of $\mathrm{D}^b(\mathbf{A})$ naturally associated to E (namely, $E^i := A_{-i}$ for $-m \leq i \leq 0$ and otherwise $E^i := 0$, with differential $d_{-i}: E^i \rightarrow E^{i+1}$ for $-m \leq i < 0$).

3.2. The criterion: extension to a subcategory. Looking carefully at the proof of [11, Prop. 3.7], one sees that the notion of ample sequence can be replaced there by the one of almost ample set. In particular, if \mathbf{T} is a triangulated category and \mathbf{A} is an abelian category, we can deal with exact functors $F: \mathrm{D}^b(\mathbf{A}) \rightarrow \mathbf{T}$ satisfying the following condition:

$$(\diamond) \quad \mathrm{Hom}_{\mathbf{T}}(F(A), F(B)[k]) = 0, \text{ for any } A, B \in \mathbf{A} \text{ and any } k < 0.$$

Hence one can prove the following result.

Proposition 3.3. *Let \mathbf{T} be a triangulated category and let \mathbf{A} be an abelian category of finite homological dimension. Assume that $\{P_i\}_{i \in I} \subseteq \mathbf{A}$ is an almost ample set and denote by \mathbf{C} the corresponding full subcategory. Let $F_1, F_2: \mathrm{D}^b(\mathbf{A}) \rightarrow \mathbf{T}$ be exact functors and let $f: F_1|_{\mathbf{C}} \xrightarrow{\sim} F_2|_{\mathbf{C}}$ be an isomorphism of functors. Assume moreover the following:*

- (i) *the functor F_1 satisfies (\diamond) ;*
- (ii) *F_1 has a left adjoint.*

Then there exists an isomorphism of exact functors $g: F_1 \xrightarrow{\sim} F_2$ extending f .

In the rest of this paper we would like to apply Proposition 3.3 but, unfortunately, in the supported case a functor $F: \mathrm{D}_{Z_1}^b(X_1) \rightarrow \mathrm{D}_{Z_2}^b(X_2)$ may not have left or right adjoint. Thus we are going to prove a more general result (Proposition 3.7, whose proof is however much inspired by those of [30, Prop. 2.16] and [11, Prop. 3.7]), from which Proposition 3.3 will follow easily (see the end of Section 3.3). To this purpose, we first introduce the categorical setting which will be used in the rest of Section 3.

Indeed, to weaken condition (\diamond) , let \mathbf{E} be a full exact subcategory of an abelian category \mathbf{A} satisfying the following conditions:

- (E1) A morphism in \mathbf{E} is an admissible epimorphism if and only if it is an epimorphism in \mathbf{A} ;
- (E2) There is a set $\{P_i\}_{i \in I} \subseteq \mathbf{E}$ which satisfies property (1) of Definition 2.9 as objects of \mathbf{A} ;
- (E3) For all $A \in D^b(\mathbf{E}) \cap \mathbf{A}$, there exists an integer $N(A)$ such that $\mathrm{Hom}_{D^b(\mathbf{A})}(A, B[i]) = 0$, for every $i > N(A)$ and every $B \in D^b(\mathbf{E}) \cap \mathbf{A}$.

The reader who is not familiar with the language of exact categories can have a look at [21] (where admissible epimorphisms are called deflations).

Remark 3.4. Under conditions (E1) and (E2), $D^b(\mathbf{E})$ can be identified with a full subcategory of $D^b(\mathbf{A})$ by [21, Thm. 12.1] and [20, Sect. 4.1] (or rather its dual version). Indeed, notice that, by (1) of Definition 2.9, for each object A of \mathbf{A} there is an object E in \mathbf{E} with an epimorphism $E \rightarrow A$ in \mathbf{A} .

For \mathbf{E} and \mathbf{A} satisfying (E1) and (E2), we will consider exact functors $F: D^b(\mathbf{E}) \rightarrow \mathbf{T}$ (for \mathbf{T} a triangulated category) such that

- (1) $\mathrm{Hom}(F(A), F(B)[k]) = 0$, for any $A, B \in D^b(\mathbf{E}) \cap \mathbf{A}$ and any integer $k < 0$;
 - (2) For all $C \in D^b(\mathbf{E})$ with trivial cohomologies in positive degrees, there is $i \in I$ such
- (Δ) that

$$\mathrm{Hom}(F(C), F(P_i)) = 0$$

and i satisfies property (1) of Definition 2.9 for $H^0(C)$.

In order to state our first extension result, we need some more notation. Let \mathbf{C} be the full subcategory of \mathbf{A} with objects $\{P_i\}_{i \in I}$ and set \mathbf{D}_0 to be the strictly full subcategory of $D^b(\mathbf{E})$ whose objects are isomorphic to shifts of objects of \mathbf{A} . Recall that a full subcategory is strictly full if it is closed under isomorphisms.

Proposition 3.5. *Let \mathbf{T} be a triangulated category and let \mathbf{E} be a full exact subcategory of an abelian category \mathbf{A} satisfying (E1), (E2) and (E3). Let $F_1, F_2: D^b(\mathbf{E}) \rightarrow \mathbf{T}$ be exact functors with a natural transformation $f: F_1|_{\mathbf{C}} \rightarrow F_2|_{\mathbf{C}}$. Assume moreover the following:*

- (i) F_1 and F_2 both satisfy condition (1) of (Δ);
- (ii) For any $A, B \in D^b(\mathbf{E}) \cap \mathbf{A}$ and any integer $k < 0$

$$\mathrm{Hom}(F_1(A), F_2(B)[k]) = 0.$$

Then there exists a unique natural transformation compatible with shifts $f_0: F_1|_{\mathbf{D}_0} \rightarrow F_2|_{\mathbf{D}_0}$ extending f .

Proof. For any $i \in I$, let $f_i := f(P_i): F_1(P_i) \rightarrow F_2(P_i)$. We also set $\mathbf{F} := D^b(\mathbf{E}) \cap \mathbf{A}$.

The first key step consists in showing that f extends uniquely to a natural transformation $F_1|_{\mathbf{F}} \rightarrow F_2|_{\mathbf{F}}$. To this purpose, one starts with $A \in \mathbf{F}$ and takes a(n infinite) resolution

$$(3.3) \quad \dots \rightarrow P_{i_j}^{\oplus k_j} \xrightarrow{d_j} P_{i_{j-1}}^{\oplus k_{j-1}} \xrightarrow{d_{j-1}} \dots \xrightarrow{d_1} P_{i_0}^{\oplus k_0} \xrightarrow{d_0} A \rightarrow 0,$$

where $i_j \in I$ and $k_j \in \mathbb{N}$ for every $j \in \mathbb{N}$. To prove the existence of such a resolution, one argues as follows. By (1) of Definition 2.9, there exist P_{i_0} , k_0 and d_0 as in (3.3). As $K_0 := \ker d_0$ is again an object of \mathbf{A} , we can apply the same argument to it, getting an epimorphism $d_1: P_{i_1}^{\oplus k_1} \rightarrow K_0$. Then one proceeds by induction.

Let $N(A)$ be as in (E3), fix $m > N(A)$ and consider the bounded complex

$$R_m := \{P_{i_m}^{\oplus k_m} \xrightarrow{d_m} P_{i_{m-1}}^{\oplus k_{m-1}} \xrightarrow{d_{m-1}} \dots \xrightarrow{d_1} P_{i_0}^{\oplus k_0}\}.$$

We can think of R_m as an object in $D^b(\mathbf{E})$ which is then a (unique up to isomorphism) convolution of R_m itself. Moreover, if we set $K_m := \ker d_m \in \mathbf{A}$, the object R_m in $D^b(\mathbf{E})$ sits in a distinguished triangle

$$K_m[m] \longrightarrow R_m \longrightarrow A \longrightarrow K_m[m+1]$$

in $D^b(\mathbf{E})$, and so $K_m \in \mathbf{F}$. Observe that, due to the choice of m , we have

$$\mathrm{Hom}_{D^b(\mathbf{A})}(A, K_m[m+1]) \cong \mathrm{Hom}_{D^b(\mathbf{A})}(A, K_m[m]) \cong 0,$$

implying $R_m \cong K_m[m] \oplus A$. Hence, we conclude that a (unique up to isomorphism) convolution of R_m is $(d_0, 0): P_{i_0}^{\oplus k_0} \rightarrow A \oplus K_m[m]$,

Hence for $q \in \{1, 2\}$ the complex

$$F_q(R_m) := \{F_q(P_{i_m}^{\oplus k_m}) \xrightarrow{F_q(d_m)} F_q(P_{i_{m-1}}^{\oplus k_{m-1}}) \xrightarrow{F_q(d_{m-1})} \dots \xrightarrow{F_q(d_1)} F_q(P_{i_0}^{\oplus k_0})\}$$

admits a convolution $(F_q(d_0), 0): F_q(P_{i_0}^{\oplus k_0}) \rightarrow F_q(A \oplus K_m[m])$. Lemma 3.1 and condition (i) ensure that such a convolution is unique up to isomorphism. Moreover, by (ii),

$$\mathrm{Hom}_{\mathbf{T}}(F_1(P_{i_j}), F_2(P_{i_k})[r]) \cong \mathrm{Hom}_{\mathbf{T}}(F_1(P_{i_l}), F_2(A)[r]) \cong 0$$

for any $i_j, i_k, i_l \in \{i_0, \dots, i_m\}$ and $r < 0$. Hence we can apply Lemma 3.2 getting a unique morphism $f_A: F_1(A) \rightarrow F_2(A)$ making the following diagram commutative:

$$\begin{array}{ccccccccccc} F_1(P_{i_m}^{\oplus k_m}) & \xrightarrow{F_1(d_m)} & F_1(P_{i_{m-1}}^{\oplus k_{m-1}}) & \xrightarrow{F_1(d_{m-1})} & \dots & \xrightarrow{F_1(d_1)} & F_1(P_{i_0}^{\oplus k_0}) & \xrightarrow{F_1(d_0)} & F_1(A) & & \\ \downarrow f_{i_m}^{\oplus k_m} & & \downarrow f_{i_{m-1}}^{\oplus k_{m-1}} & & & & \downarrow f_{i_0}^{\oplus k_0} & & \downarrow f_A & & \\ F_2(P_{i_m}^{\oplus k_m}) & \xrightarrow{F_2(d_m)} & F_2(P_{i_{m-1}}^{\oplus k_{m-1}}) & \xrightarrow{F_2(d_{m-1})} & \dots & \xrightarrow{F_2(d_1)} & F_2(P_{i_0}^{\oplus k_0}) & \xrightarrow{F_2(d_0)} & F_2(A). & & \end{array}$$

By Lemma 3.2, the definition of f_A does not depend on the choice of m . In other words, if we choose a different $m' > N(A)$ and truncate (3.3) in position m' , the bounded complexes $F_q(R_{m'})$ give rise to the same morphism f_A .

To show that the definition of f_A does not depend on the choice of the resolution (3.3), consider another resolution of A

$$(3.4) \quad \dots \rightarrow P_{i'_j}^{\oplus k'_j} \xrightarrow{d'_j} P_{i'_{j-1}}^{\oplus k'_{j-1}} \xrightarrow{d'_{j-1}} \dots \xrightarrow{d'_1} P_{i'_0}^{\oplus k'_0} \xrightarrow{d'_0} A \rightarrow 0.$$

and denote by $f'_A: F_1(A) \rightarrow F_2(A)$ the induced morphism. In order to see that $f_A = f'_A$, we start by proving that there exist a third resolution

$$(3.5) \quad \dots \rightarrow P_{i''_j}^{\oplus k''_j} \xrightarrow{d''_j} P_{i''_{j-1}}^{\oplus k''_{j-1}} \xrightarrow{d''_{j-1}} \dots \xrightarrow{d''_1} P_{i''_0}^{\oplus k''_0} \xrightarrow{d''_0} A \rightarrow 0$$

and morphisms $s_j: P_{i_j}^{\oplus k_j} \rightarrow P_{i_j}^{\oplus k'_j}$ and $t_j: P_{i_j}^{\oplus k''_j} \rightarrow P_{i_j}^{\oplus k'_j}$, for any $j \geq 0$, fitting into the following commutative diagram:

$$(3.6) \quad \begin{array}{ccccccc} \cdots & \xrightarrow{d'_{j+1}} & P_{i_j}^{\oplus k'_j} & \xrightarrow{d'_j} & P_{i_{j-1}}^{\oplus k'_{j-1}} & \xrightarrow{d'_{j-1}} & \cdots \xrightarrow{d'_1} P_{i_0}^{\oplus k'_0} \\ & & \uparrow t_j & & \uparrow t_{j-1} & & \uparrow t_0 \\ \cdots & \xrightarrow{d''_{j+1}} & P_{i_j}^{\oplus k''_j} & \xrightarrow{d''_j} & P_{i_{j-1}}^{\oplus k''_{j-1}} & \xrightarrow{d''_{j-1}} & \cdots \xrightarrow{d''_1} P_{i_0}^{\oplus k''_0} \\ & & \downarrow s_j & & \downarrow s_{j-1} & & \downarrow s_0 \\ \cdots & \xrightarrow{d_{j+1}} & P_{i_j}^{\oplus k_j} & \xrightarrow{d_j} & P_{i_{j-1}}^{\oplus k_{j-1}} & \xrightarrow{d_{j-1}} & \cdots \xrightarrow{d_1} P_{i_0}^{\oplus k_0} \end{array} \begin{array}{l} \searrow d'_0 \\ \nearrow d_0 \end{array} \rightarrow A.$$

In fact, we can actually argue as follows. Let $F''_0 := P_{i_0}^{\oplus k'_0} \times_A P_{i_0}^{\oplus k_0}$. By (1) of Definition 2.9, there exists an epimorphism $d''_0: P_{i_0}^{\oplus k''_0} \rightarrow F''_0 \rightarrow A$ with maps s_0 and t_0 sitting, by definition, in the commutative diagram

$$\begin{array}{ccc} P_{i_0}^{\oplus k'_0} & & \\ t_0 \uparrow & \searrow d'_0 & \\ P_{i_0}^{\oplus k''_0} & \xrightarrow{d''_0} & A \\ s_0 \downarrow & \nearrow d_0 & \\ P_{i_0}^{\oplus k_0} & & \end{array}$$

Let us now explain how we deal with the inductive step. Assume we have constructed the following commutative diagram

$$\begin{array}{ccccccc} \cdots & \xrightarrow{d'_{j+1}} & P_{i_j}^{\oplus k'_j} & \xrightarrow{d'_j} & P_{i_{j-1}}^{\oplus k'_{j-1}} & \xrightarrow{d'_{j-1}} & \cdots \xrightarrow{d'_1} P_{i_0}^{\oplus k'_0} \\ & & \uparrow t_{j-1} & & \uparrow t_0 & & \uparrow t_0 \\ & & P_{i_{j-1}}^{\oplus k''_{j-1}} & \xrightarrow{d''_{j-1}} & \cdots \xrightarrow{d''_1} & P_{i_0}^{\oplus k''_0} & \xrightarrow{d''_0} A \\ & & \downarrow s_{j-1} & & \downarrow s_0 & & \downarrow s_0 \\ \cdots & \xrightarrow{d_{j+1}} & P_{i_j}^{\oplus k_j} & \xrightarrow{d_j} & P_{i_{j-1}}^{\oplus k_{j-1}} & \xrightarrow{d_{j-1}} & \cdots \xrightarrow{d_1} P_{i_0}^{\oplus k_0} \end{array} \begin{array}{l} \searrow d'_0 \\ \nearrow d_0 \end{array} \rightarrow A.$$

We can define $P_{i_j}^{\oplus k''_j}$, d''_j , s_j and t_j as follows. Set $K'_{j-1} := \ker d'_{j-1}$, $K''_{j-1} := \ker d''_{j-1}$ and $K_{j-1} := \ker d_{j-1}$. By definition, we have $d_{j-1}(s_{j-1}(K''_{j-1})) = d'_{j-1}(t_{j-1}(K''_{j-1})) = 0$ and thus s_{j-1} and t_{j-1} induce two morphisms $r'_{j-1}: K''_{j-1} \rightarrow K'_{j-1}$ and $r_{j-1}: K''_{j-1} \rightarrow K_{j-1}$. Consider the fiber products

$$F'_{j-1} := P_{i_j}^{\oplus k'_j} \times_{K'_{j-1}} K''_{j-1} \quad F_{j-1} := P_{i_j}^{\oplus k_j} \times_{K_{j-1}} K''_{j-1}$$

along the maps d'_j , r'_{j-1} and d_j , r_{j-1} respectively. Since d'_j and d_j are epimorphisms onto K'_{j-1} and K_{j-1} respectively, we get epimorphisms $F'_{j-1} \rightarrow K''_{j-1}$ and $F_{j-1} \rightarrow K''_{j-1}$. Take the fiber

product $F''_{j-1} := F'_{j-1} \times_{K''_{j-1}} F_{j-1}$ which then comes with an epimorphism onto K''_{j-1} . By (1) of Definition 2.9, we get an epimorphism $d''_j: P_{i''_j}^{\oplus k''_j} \twoheadrightarrow F''_{j-1} \twoheadrightarrow K''_{j-1}$ with maps s_j and t_j making (3.6) commutative.

Denoting by $f''_A: F_1(A) \rightarrow F_2(A)$ the morphism constructed using (3.5), we get a diagram

$$\begin{array}{ccccc}
F_1(P_{i''_0}^{\oplus k''_0}) & \xrightarrow{F_1(d''_0)} & & & F_1(A) \\
\downarrow f_{i''_0}^{\oplus k''_0} & \searrow F_1(s_0) & & & \downarrow \text{id} \\
& & F_1(P_{i_0}^{\oplus k_0}) & \xrightarrow{F_1(d_0)} & F_1(A) \\
& & \downarrow f_{i_0}^{\oplus k_0} & & \downarrow f_A \quad \star \\
& & F_2(P_{i_0}^{\oplus k_0}) & \xrightarrow{F_2(d_0)} & F_2(A) \\
& \nearrow F_2(s_0) & & & \uparrow \text{id} \\
F_2(P_{i''_0}^{\oplus k''_0}) & \xrightarrow{F_2(d''_0)} & & & F_2(A) \\
& & & & \downarrow f''_A
\end{array}$$

where all squares and trapezoids but \star are commutative. Due to hypothesis (ii) and Lemma 3.2 there exists a unique morphism $F_1(A) \rightarrow F_2(A)$ making the following diagram commutative:

$$\begin{array}{ccc}
F_1(P_{i''_0}^{\oplus k''_0}) & \xrightarrow{F_1(d''_0)} & F_1(A) \\
F_2(s_0) \circ f_{i''_0}^{\oplus k''_0} \downarrow & & \downarrow \\
F_2(P_{i_0}^{\oplus k_0}) & \xrightarrow{F_2(d_0)} & F_2(A).
\end{array}$$

Since $F_2(s_0) \circ f_{i''_0}^{\oplus k''_0} = f_{i_0}^{\oplus k_0} \circ F_1(s_0)$ by the naturality of f on \mathbf{C} , both f_A and f''_A have this property and then they coincide. Similarly one can prove that $f''_A = f'_A$.

Therefore $f_A = f'_A$, and so $\tilde{f}(A) := f_A: F_1(A) \rightarrow F_2(A)$ is well defined for every $A \in \mathbf{F} = \mathbf{D}^b(\mathbf{E}) \cap \mathbf{A}$. It is also easy to see that $\tilde{f}: \mathbf{F}_1|_{\mathbf{F}} \rightarrow \mathbf{F}_2|_{\mathbf{F}}$ is a natural transformation. Indeed, let $u: A \rightarrow B$ be a morphism of \mathbf{F} . Consider a resolution of B

$$\dots \rightarrow P_{l_j}^{\oplus h_j} \xrightarrow{e_j} P_{l_{j-1}}^{\oplus h_{j-1}} \xrightarrow{e_{j-1}} \dots \xrightarrow{e_1} P_{l_0}^{\oplus h_0} \xrightarrow{e_0} B \rightarrow 0,$$

where $l_j \in I$ and $h_j \in \mathbb{N}$ for every $j \in \mathbb{N}$. We can find a resolution of A

$$\dots \rightarrow P_{i_j}^{\oplus k_j} \xrightarrow{d_j} P_{i_{j-1}}^{\oplus k_{j-1}} \xrightarrow{d_{j-1}} \dots \xrightarrow{d_1} P_{i_0}^{\oplus k_0} \xrightarrow{d_0} A \rightarrow 0$$

and morphisms $g_j: P_{i_j}^{\oplus k_j} \rightarrow P_{l_j}^{\oplus h_j}$ defining a morphism of complexes compatible with u . Indeed, set $E_0 := A \times_B P_{l_0}^{\oplus h_0}$ to be the fiber product along the morphisms u and e_0 . By (1) of Definition 2.9, we get an epimorphism $P_{i_0}^{\oplus k_0} \twoheadrightarrow E_0$ providing, by composition, the desired morphisms d_0 and $g_0: P_{i_0}^{\oplus k_0} \rightarrow P_{l_0}^{\oplus h_0}$. Then we proceed by induction as in (3.6).

We can now consider the diagram

$$\begin{array}{ccccc}
F_1(P_{i_0}^{\oplus k_0}) & \xrightarrow{F_1(d_0)} & & \xrightarrow{F_1(d_0)} & F_1(A) \\
\downarrow F_1(g_0) & \searrow f_{i_0}^{\oplus k_0} & & \swarrow f_A & \downarrow F_1(u) \\
& & F_2(P_{i_0}^{\oplus k_0}) & \xrightarrow{F_2(d_0)} & F_2(A) \\
& & \downarrow F_2(g_0) & & \downarrow F_2(u) \\
& & F_2(P_{l_0}^{\oplus h_0}) & \xrightarrow{F_2(e_0)} & F_2(B) \\
& \nearrow f_{i_0}^{\oplus h_0} & & \nwarrow f_B & \\
F_1(P_{l_0}^{\oplus h_0}) & \xrightarrow{F_1(e_0)} & & \xrightarrow{F_1(e_0)} & F_1(B)
\end{array}$$

where all squares and trapezoids but \star are commutative. Using the same argument as above, we can take $m > N(A), N(B)$ and truncate the resolutions of A and B at step m . Then, applying (ii) and Lemma 3.2, we see that there is a unique morphism $F_1(A) \rightarrow F_2(B)$ completing the following diagram to a commutative square

$$\begin{array}{ccc}
F_1(P_{i_0}^{\oplus k_0}) & \xrightarrow{F_1(d_0)} & F_1(A) \\
F_2(g_0) \circ f_{i_0}^{\oplus k_0} \downarrow & & \downarrow \\
F_2(P_{l_0}^{\oplus h_0}) & \xrightarrow{F_2(e_0)} & F_2(B).
\end{array}$$

Since $F_2(g_0) \circ f_{i_0}^{\oplus k_0} = f_{l_0}^{\oplus h_0} \circ F_1(g_0)$, both $F_2(u) \circ f_A$ and $f_B \circ F_1(u)$ have this property. It follows that $F_2(u) \circ \tilde{f}(A) = \tilde{f}(B) \circ F_1(u)$, thus proving that $\tilde{f}: F_1|_{\mathbf{F}} \rightarrow F_2|_{\mathbf{F}}$ is a natural transformation. It is clear by construction that $\tilde{f}|_{\mathbf{C}} = f$ and that \tilde{f} is unique with this property.

Since the objects of \mathbf{D}_0 are precisely (up to isomorphism) shifts of objects of \mathbf{F} , we just need to define $f_0(A[k])$ for $A \in \mathbf{F}$ and $k \in \mathbb{Z}$. Of course, we must set $f_0(A[k]) := \tilde{f}(A)[k]$, and we have just to show that $f_0(B[k]) \circ F_1(u) = F_2(u) \circ f_0(A)$ for all objects $A, B \in \mathbf{F}$ and every $u \in \text{Hom}(A, B[k])$. Notice that, for simplicity, we are not making explicit the natural isomorphism $F_i(A[k]) \cong F_i(A)[k]$, for $i = 1, 2$.

Now, there is nothing to prove if $k < 0$ (because then $u = 0$) or $k = 0$ (because we have already seen that \tilde{f} is a natural transformation), so we assume $k > 0$. Actually we can reduce to the case $k = 1$, thanks to the fact that one can always factor u as $u = u_k \circ \cdots \circ u_1$, where (for $j = 1, \dots, k$) $u_j \in \text{Hom}(C_{j-1}[j-1], C_j[j])$ and $C_0 = A, \dots, C_k = B$ are objects of \mathbf{F} (see Step 4 in the proof of [28, Prop. B.1]). Now, completing u to a distinguished triangle $B \rightarrow C \xrightarrow{v} A \xrightarrow{u} B[1]$, C is again an object of \mathbf{F} . Then by axiom (TR3) there exists a morphism $h: F_1(A) \rightarrow F_2(A)$ such that the diagram

$$\begin{array}{ccccccc}
F_1(B) & \longrightarrow & F_1(C) & \xrightarrow{F_1(v)} & F_1(A) & \xrightarrow{F_1(u)} & F_1(B[1]) \\
\downarrow \tilde{f}(B) & & \downarrow \tilde{f}(C) & & \downarrow h & & \downarrow f_0(B[1]) \\
F_2(B) & \longrightarrow & F_2(C) & \xrightarrow{F_2(v)} & F_2(A) & \xrightarrow{F_2(u)} & F_2(B[1])
\end{array}$$

commutes. Since $\text{Hom}(\mathbf{F}_1(B[1]), \mathbf{F}_2(A)) = 0$ by hypothesis, h is the unique morphism such that $h \circ \mathbf{F}_1(v) = \mathbf{F}_2(v) \circ \tilde{f}(C)$. Hence $h = \tilde{f}(A) = f_0(A)$, and we conclude that $f_0(B[1]) \circ \mathbf{F}_1(u) = \mathbf{F}_2(u) \circ f_0(A)$. \square

Let us specialize to the case of isomorphisms.

Corollary 3.6. *With the same hypotheses on \mathbf{E} , \mathbf{A} and \mathbf{T} , let $\mathbf{F}_1, \mathbf{F}_2: \mathbf{D}^b(\mathbf{E}) \rightarrow \mathbf{T}$ be exact functors and let $f: \mathbf{F}_1|_{\mathbf{C}} \xrightarrow{\sim} \mathbf{F}_2|_{\mathbf{C}}$ be an isomorphism of functors. Assume that \mathbf{F}_1 satisfies condition (1) of (Δ) . Then there exists a unique isomorphism compatible with shifts $f_0: \mathbf{F}_1|_{\mathbf{D}_0} \xrightarrow{\sim} \mathbf{F}_2|_{\mathbf{D}_0}$ extending f .*

Proof. Since f is an isomorphism, we can use Lemma 3.1 in the above argument to show that there is an isomorphism $\mathbf{F}_1(A) \cong \mathbf{F}_2(A)$, for all $A \in \mathbf{D}^b(\mathbf{E}) \cap \mathbf{A}$. Thus (ii) in Proposition 3.5 follows from (1) in (Δ) . Therefore, by the above proposition, there is a unique natural transformation compatible with shifts $f_0: \mathbf{F}_1 \rightarrow \mathbf{F}_2$ extending f . Uniqueness, applied also to f^{-1} , $f \circ f^{-1}$ and $f^{-1} \circ f$, immediately implies that f_0 is an isomorphism \square

3.3. The criterion: extension to the whole derived category. In order to extend the natural transformation f of Proposition 3.5 to $\mathbf{D}^b(\mathbf{E})$, the exact functors have to satisfy one more assumption.

Proposition 3.7. *Let \mathbf{T} be a triangulated category and let \mathbf{E} be a full exact subcategory of an abelian category \mathbf{A} satisfying (E1), (E2) and (E3). Let $\mathbf{F}_1, \mathbf{F}_2: \mathbf{D}^b(\mathbf{E}) \rightarrow \mathbf{T}$ be exact functors with a natural transformation $f: \mathbf{F}_1|_{\mathbf{C}} \rightarrow \mathbf{F}_2|_{\mathbf{C}}$. Assume moreover the following:*

- (i) \mathbf{F}_1 and \mathbf{F}_2 both satisfy condition (1) of (Δ) ;
- (ii) For any $A, B \in \mathbf{D}^b(\mathbf{E}) \cap \mathbf{A}$ and any integer $k < 0$

$$\text{Hom}(\mathbf{F}_1(A), \mathbf{F}_2(B)[k]) = 0.$$

- (iii) for all $C \in \mathbf{D}^b(\mathbf{E})$ with trivial cohomologies in positive degrees, there is $i \in I$ such that

$$\text{Hom}(\mathbf{F}_1(C), \mathbf{F}_2(P_i)) = 0$$

and i satisfies condition (1) of Definition 2.9 for $H^0(C)$.

Then there exists a unique natural transformation of exact functors $g: \mathbf{F}_1 \rightarrow \mathbf{F}_2$ extending f .

Proof. For $n \in \mathbb{N}$, denote by \mathbf{D}_n the (strictly) full subcategory of $\mathbf{D}^b(\mathbf{E})$ with objects the complexes A with the following property: there exists $a \in \mathbb{Z}$ such that $H^p(A) = 0$ for $p < a$ or $p > a + n$. We are going to prove by induction on n that f extends uniquely to a natural transformation compatible with shifts $f_n: \mathbf{F}_1|_{\mathbf{D}_n} \rightarrow \mathbf{F}_2|_{\mathbf{D}_n}$. Once we do this, it is obvious that for every object A of $\mathbf{D}^b(\mathbf{E})$ we can define $g(A) := f_n(A)$ if $A \in \mathbf{D}_n$, and that g is then the unique required extension of f .

The case $n = 0$ having already been proved in Proposition 3.5, we come to the inductive step from $n - 1$ to $n > 0$. For every object $A \in \mathbf{D}_n$ we need to define $f_n(A): \mathbf{F}_1(A) \rightarrow \mathbf{F}_2(A)$. To this purpose, we can assume that $H^p(A) = 0$ for $p < -n$ or $p > 0$. If $A = \{\dots \rightarrow A^0 \xrightarrow{d^0} A^1 \rightarrow \dots\}$, let

$$s: P_i^{\oplus k} \twoheadrightarrow \ker d^0$$

(for some $i \in I$ and $k \in \mathbb{N}$) be an epimorphism such that $\text{Hom}(\mathbf{F}_1(A), \mathbf{F}_2(P_i)) = 0$. Notice that s can be found as follows: after choosing an epimorphism $P_j^{\oplus l} \twoheadrightarrow \ker d^0$ (with $j \in I$ and $l \in \mathbb{N}$), take $i \in I$ which satisfies condition (iii) for $A \oplus P_j^{\oplus l}$ (so that, in particular, there is an epimorphism $P_i^{\oplus k} \twoheadrightarrow P_j^{\oplus l}$), and define s to be the composition $P_i^{\oplus k} \twoheadrightarrow P_j^{\oplus l} \twoheadrightarrow \ker d^0$. Denoting by $t: P_i^{\oplus k} \rightarrow A$ the composition of s with the natural morphism $\ker d^0 \rightarrow A$, it is then clear that $H^0(t): P_i^{\oplus k} \rightarrow H^0(A)$ is an epimorphism. It follows that we have a distinguished triangle

$$(3.7) \quad C[-1] \rightarrow P_i^{\oplus k} \xrightarrow{t} A \xrightarrow{t_1} C$$

with $C \in \mathbf{D}_{n-1}$. Hence, by the inductive hypothesis and using axiom (TR3), we obtain a commutative diagram whose rows are distinguished triangles

$$\begin{array}{ccccccc} \mathbf{F}_1(C)[-1] & \longrightarrow & \mathbf{F}_1(P_i^{\oplus k}) & \xrightarrow{\mathbf{F}_1(t)} & \mathbf{F}_1(A) & \xrightarrow{\mathbf{F}_1(t_1)} & \mathbf{F}_1(C) \\ \downarrow f_{n-1}(C)[-1] & & \downarrow f_{n-1}(P_i^{\oplus k}) & & \downarrow f_A & & \downarrow f_{n-1}(C) \\ \mathbf{F}_2(C)[-1] & \longrightarrow & \mathbf{F}_2(P_i^{\oplus k}) & \xrightarrow{\mathbf{F}_2(t)} & \mathbf{F}_2(A) & \xrightarrow{\mathbf{F}_2(t_1)} & \mathbf{F}_2(C) \end{array}$$

for some $f_A: \mathbf{F}_1(A) \rightarrow \mathbf{F}_2(A)$. Observe that, since $\text{Hom}(\mathbf{F}_1(A), \mathbf{F}_2(P_i^{\oplus k})) = 0$ by assumption, f_A is the unique morphism such that the square on the right commutes.

In order to prove that f_A does not depend on the choice of s , assume that $s': P_{i'}^{\oplus k'} \twoheadrightarrow \ker d^0$ is another epimorphism such that $\text{Hom}(\mathbf{F}_1(A), \mathbf{F}_2(P_{i'})) = 0$, and thus inducing another morphism $f'_A: \mathbf{F}_1(A) \rightarrow \mathbf{F}_2(A)$. We claim that we can find a third epimorphism $s'': P_{i''}^{\oplus k''} \twoheadrightarrow \ker d^0$ such that $\text{Hom}(\mathbf{F}_1(A), \mathbf{F}_2(P_{i''})) = 0$ (inducing $f''_A: \mathbf{F}_1(A) \rightarrow \mathbf{F}_2(A)$) and fitting into a commutative diagram

$$\begin{array}{ccc} P_{i''}^{\oplus k''} & \xrightarrow{w} & P_i^{\oplus k} \\ \downarrow w' & \searrow s'' & \downarrow s \\ P_{i'}^{\oplus k'} & \xrightarrow{s'} & \ker d^0 \end{array}$$

This can be easily seen if one takes $i'' \in I$ satisfying condition (iii) for $A \oplus P_j^{\oplus l}$, where $j \in I$ and $l \in \mathbb{N}$ are such that there exists an epimorphism $P_j^{\oplus l} \twoheadrightarrow P_i^{\oplus k} \times_{\ker d^0} P_{i'}^{\oplus k'}$ (then s'' is an epimorphism because the natural map $P_i^{\oplus k} \times_{\ker d^0} P_{i'}^{\oplus k'} \twoheadrightarrow \ker d^0$ is an epimorphism). Observing that the morphisms $t': P_{i'}^{\oplus k'} \rightarrow A$ and $t'': P_{i''}^{\oplus k''} \rightarrow A$ (induced, respectively, by s' and s'') obviously satisfy $t \circ w = t'' = t' \circ w'$, by axiom (TR3) there is a commutative diagram whose rows are distinguished triangles

$$\begin{array}{ccccccc} P_{i''}^{\oplus k''} & \xrightarrow{t''} & A & \xrightarrow{t'_1} & C'' & \longrightarrow & P_{i''}^{\oplus k''}[1] \\ \downarrow w & & \downarrow \text{id} & & \downarrow v & & \downarrow w[1] \\ P_i^{\oplus k} & \xrightarrow{t} & A & \xrightarrow{t_1} & C & \longrightarrow & P_i^{\oplus k}[1] \end{array}$$

for some $v: C'' \rightarrow C$. As the diagram

$$\begin{array}{ccccc} F_1(A) & \xrightarrow{F_1(t_1'')} & F_1(C'') & \xrightarrow{F_1(v)} & F_1(C) \\ \downarrow f_A'' & & \downarrow f_{n-1}(C'') & & \downarrow f_{n-1}(C) \\ F_2(A) & \xrightarrow{F_2(t_1'')} & F_2(C'') & \xrightarrow{F_2(v)} & F_2(C) \end{array}$$

commutes (the square on the left by definition of f_A'' , the square on the right because f_{n-1} is a natural transformation by induction) and since $v \circ t_1'' = t_1$, we obtain

$$f_{n-1}(C) \circ F_1(t_1) = f_{n-1}(C) \circ F_1(v) \circ F_1(t_1'') = F_2(v) \circ F_2(t_1'') \circ f_A'' = F_2(t_1) \circ f_A''.$$

On the other hand, f_A is the only morphism with the property that $f_{n-1}(C) \circ F_1(t_1) = F_2(t_1) \circ f_A$. It follows that $f_A = f_A''$ and similarly $f_A' = f_A''$, thereby proving that $f_A = f_A'$. Therefore we can set $f_n(A) := f_A$, and more generally $f_n(A[k]) := f_A[k]$ for every integer k , thus defining f_n on every object of \mathbf{D}_n .

To conclude the inductive step it is enough to show that f_n is a natural transformation, because then it is clear by definition that f_n is compatible with shifts, that $f_n|_C = f_{n-1}|_C = f$ (actually also $f_n|_{\mathbf{D}_{n-1}} = f_{n-1}$) and that f_n is unique with these properties. So we have to prove that

$$(3.8) \quad f_n(B) \circ F_1(u) = F_2(u) \circ f_n(A)$$

for every morphism $u: A \rightarrow B$ of \mathbf{D}_n . If (3.8) holds for a morphism u of \mathbf{D}_n , we say that f_n is *compatible* with u .

Recall that in $D^b(\mathbf{E})$ we can write $u = w_1 \circ w_2^{-1}$, where w_1 and w_2 are (represented by) morphisms of complexes and w_2 is a quasi-isomorphism (hence w_1 and w_2 are again in \mathbf{D}_n). Thus f_n is compatible with u if it is compatible both with w_2^{-1} (or, equivalently, with w_2) and with w_1 . In other words, it is harmless to assume directly that u is a morphism of complexes, denoted by $A = \{\dots \rightarrow A^0 \xrightarrow{d^0} A^1 \rightarrow \dots\}$ and $B = \{\dots \rightarrow B^0 \xrightarrow{e^0} B^1 \rightarrow \dots\}$. We can also assume that, as before, $H^p(A) = 0$ for $p < -n$ or $p > 0$. Moreover, we denote by c the greatest integer such that $H^c(B) \neq 0$ (of course, if $B \cong 0$ there is nothing to prove). Now our aim is to show that the problem of verifying (3.8) can be reduced to a similar problem with another ‘‘simpler’’ morphism in place of u . To this purpose we distinguish two cases according to the value of c .

Case 1: $c < 0$. Choose $j \in I$ which satisfies (1) of Definition 2.9 for $K := \ker d^0 \times_{\ker e^0} B^{-1}$, let $P_j^{\oplus l} \twoheadrightarrow K$ be an epimorphism, and take $i \in I$ satisfying condition (iii) for $A \oplus P_j^{\oplus l}$. Then, reasoning as before, we get an epimorphism $s: P_i^{\oplus k} \twoheadrightarrow \ker d^0$ (notice that $K \rightarrow \ker d^0$ is an epimorphism because $B^{-1} \rightarrow \ker e^0 = \text{im } e^{-1}$ is an epimorphism, as $c < 0$) which can be used to define f_A . Moreover, denoting by $t: P_i^{\oplus k} \rightarrow A$ the morphism (of complexes) induced by s , it is clear that $u \circ t$ is given by a morphism $w: P_i^{\oplus k} \rightarrow \ker e^0 \subseteq B^0$ which factors through the natural morphism $K \rightarrow \ker e^0$. In particular, there exists $w': P_i^{\oplus k} \rightarrow B^{-1}$ such that $w = e^{-1} \circ w'$. This proves that $u \circ t$ is homotopic to 0, whence it is 0 in $D^b(\mathbf{E})$. From this and from the distinguished triangle (3.7) it follows that $u = v \circ t_1$ for some $v: C \rightarrow B$ (with $C \in \mathbf{D}_{n-1}$). As f_n is compatible with t_1 by definition of $f_A = f_n(A)$, in order to check (3.8) it is therefore enough to show that f_n is compatible with v . Notice that, if $A \in \mathbf{D}_m$ for some $0 < m \leq n$, then $C \in \mathbf{D}_{m-1}$. On the other

hand, if $A \in \mathbf{D}_0$ (hence A is isomorphic to an object of \mathbf{F}), then $C \in \mathbf{D}_0$ and $C[-1]$ is isomorphic to an object of \mathbf{F} . So in this last case, passing from u to $v[-1]$, c increases by 1.

Case 2: $c \geq 0$. Choose an epimorphism $P_j^{\oplus l} \rightarrow \ker e^c$ (with $j \in I$ and $l \in \mathbb{N}$) and take $i \in I$ satisfying condition (iii) for $A[c] \oplus B[c] \oplus P_j^{\oplus l}$. Then, as usual, we can find an epimorphism $s': P_i^{\oplus k} \rightarrow \ker e^c$ which can be used to define $f_{B[c]}$. Denoting by $t': P_i^{\oplus k}[-c] \rightarrow B$ the morphism induced by s' , and extending it to a distinguished triangle

$$C'[-1] \rightarrow P_i^{\oplus k}[-c] \xrightarrow{t'} B \xrightarrow{t'_1} C'$$

(with $C' \in \mathbf{D}_{n-1}$), we claim that (3.8) follows once one proves that f_n is compatible with $v' := t'_1 \circ u: A \rightarrow C'$. To see this, observe that in the diagram

$$\begin{array}{ccccc} F_1(A) & \xrightarrow{F_1(u)} & F_1(B) & \xrightarrow{F_1(t'_1)} & F_1(C') \\ \downarrow f_n(A) & & \downarrow f_n(B) & & \downarrow f_n(C') \\ F_2(A) & \xrightarrow{F_2(u)} & F_2(B) & \xrightarrow{F_2(t'_1)} & F_2(C') \end{array}$$

the square on the right commutes by definition of $f_{B[c]}[-c] = f_n(B)$, whence (assuming compatibility of f_n with v')

$$F_2(t'_1) \circ (f_n(B) \circ F_1(u) - F_2(u) \circ f_n(A)) = f_n(C') \circ F_1(v') - F_2(v') \circ f_n(A) = 0.$$

It follows that $f_n(B) \circ F_1(u) - F_2(u) \circ f_n(A)$ factors through $F_2(t')$, and then it must be 0 (which means that (3.8) holds) because $\text{Hom}(F_1(A), F_2(P_i^{\oplus k}[-c])) = 0$ by the choice of i . Observe that, similarly as above, if $B \in \mathbf{D}_m$ for some $0 < m \leq n$, then $C' \in \mathbf{D}_{m-1}$, whereas, if $B \in \mathbf{D}_0$, then $C' \in \mathbf{D}_0$ and, passing from u to v' , c decreases by 1.

To finish the proof, note that, applying the above procedure, one obtains a morphism having the source in \mathbf{D}_{n-1} and the same target (in Case 1) or the same source and the target in \mathbf{D}_{n-1} (in Case 2). So it is enough to show that, repeating the procedure a sufficient number of times, one necessarily encounters both cases (because then one reduces to check compatibility of f_n with a morphism of \mathbf{D}_{n-1} , where it holds by induction). Indeed, if one always encounters Case 1 (the argument is completely similar for Case 2), then in a finite number (at most n) of steps the source becomes an object of \mathbf{D}_0 . Applying another finite number of steps, one eventually gets $c = 0$, namely Case 2. \square

In the paper we will need the following special case of the above result.

Corollary 3.8. *With the same hypotheses on \mathbf{E} , \mathbf{A} and \mathbf{T} , let $F_1, F_2: D^b(\mathbf{E}) \rightarrow \mathbf{T}$ be exact functors and let $f: F_1|_{\mathbf{C}} \xrightarrow{\sim} F_2|_{\mathbf{C}}$ be an isomorphism. Assume moreover that F_1 satisfies (Δ) . Then there exists a unique isomorphism of exact functors $g: F_1 \xrightarrow{\sim} F_2$ extending f .*

Proof. As f is an isomorphism, we can apply Corollary 3.6 so that $F_1(A) \cong F_2(A)$, for all $A \in D^b(\mathbf{E}) \cap \mathbf{A}$. Hence hypotheses (i) and (ii) in Proposition 3.7 follow from (Δ) . Analogously, for (iii) we use that $F_1(P_i) \cong F_2(P_i)$ by assumption. Thus Proposition 3.7 applies and we get a unique natural transformation of exact functors $g: F_1 \rightarrow F_2$ extending f . The fact that g is an isomorphism is again a formal consequence of uniqueness, as in the proof of Corollary 3.6. \square

In the case $\mathbf{E} = \mathbf{A}$, we are going to give a sufficient condition under which (Δ) is automatically satisfied. We leave it to the reader to formulate a similar statement which ensures that the hypotheses of Proposition 3.7 are satisfied.

Lemma 3.9. *Let $F: D^b(\mathbf{A}) \rightarrow \mathbf{T}$ be an exact functor admitting a left adjoint and satisfying (\diamond) . Assume moreover that $\{P_i\}_{i \in I}$ is an almost ample set in \mathbf{A} . Then F satisfies (Δ) as well.*

Proof. Observing that part (1) of (Δ) coincides with (\diamond) because $\mathbf{E} = \mathbf{A}$, it remains to prove part (2) of (Δ) . Denoting by $F^*: \mathbf{T} \rightarrow D^b(\mathbf{A})$ the left adjoint of F , we claim that $H^p(F^* \circ F(A)) = 0$ for any $A \in \mathbf{A}$ and for any $p > 0$. Indeed, otherwise there would exist $A \in \mathbf{A}$ and $m > 0$ with a non-zero morphism $F^* \circ F(A) \rightarrow H^m(F^* \circ F(A))[-m]$ (it is enough to let m be the largest integer such that $H^m(F^* \circ F(A)) \neq 0$). But then

$$0 \neq \mathrm{Hom}(F^* \circ F(A), H^m(F^* \circ F(A))[-m]) \cong \mathrm{Hom}(F(A), F(H^m(F^* \circ F(A))[-m]))$$

by adjunction, contradicting (\diamond) .

The above implies more generally that $H^p(F^* \circ F(C)) = 0$, for any $C \in D^b(\mathbf{A})$ having trivial cohomologies in positive degrees and any $p > 0$. To see this, we can proceed by induction on the smallest integer n such that C is an object of \mathbf{D}_n (the full subcategory of $D^b(\mathbf{E}) = D^b(\mathbf{A})$ defined at the beginning of the proof of Proposition 3.7). Indeed, we can assume without loss of generality that $H^0(C)$ is the greatest non-trivial cohomology of C . Then C is isomorphic to an object of \mathbf{A} if $n = 0$, so the statement has already been proved in this case. If $n > 0$ we have a distinguished triangle

$$C' \rightarrow C \rightarrow H^0(C) \rightarrow C'[1]$$

with C' having non-trivial cohomologies only in negative degrees and $C' \in \mathbf{D}_{n-1}$. By induction, for $p > 0$, we have $H^p(F^* \circ F(C')) = H^p(F^* \circ F(H^0(C))) = 0$. Thus $H^p(F^* \circ F(C)) = 0$.

Then for such an object C and for any $i \in I$ we have

$$\mathrm{Hom}(F(C), F(P_i)) \cong \mathrm{Hom}(F^* \circ F(C), P_i) \cong \mathrm{Hom}(H^0(F^* \circ F(C)), P_i).$$

For the last isomorphism above, take the distinguished triangle

$$C'' \rightarrow F^* \circ F(C) \rightarrow H^0(F^* \circ F(C)) \rightarrow C''[1],$$

where again, C'' has cohomologies in degrees smaller than zero. Hence

$$\mathrm{Hom}(C'', P_i) \cong \mathrm{Hom}(C''[1], P_i) \cong 0.$$

Therefore part (2) of (Δ) is satisfied if one takes $i \in I$ as in Definition 2.9 for $H^0(F^* \circ F(C)) \oplus H^0(C)$. \square

Combining the above result with Corollary 3.8 immediately gives a proof of Proposition 3.3.

3.4. The geometric case and some examples. In this section we want to clarify which abelian category \mathbf{A} and exact subcategory \mathbf{E} have to be taken in order to use the results in Section 3.3 to prove Theorems 1.1 and 1.2.

Therefore let X be a quasi-projective scheme and let Z be a projective subscheme of X . Assume further that $\mathcal{O}_{iZ} \in \mathbf{Perf}(X)$ for all $i > 0$. Set

$$\mathbf{A} := \mathbf{Coh}_Z(X) \quad \mathbf{E} := \mathbf{Perf}_Z(X) \cap \mathbf{Coh}_Z(X).$$

Proposition 3.10. *Under the above assumptions, \mathbf{E} is a full exact subcategory of \mathbf{A} , (E1), (E2) and (E3) are satisfied and $\mathbf{Perf}_Z(X) = D^b(\mathbf{E}) \subseteq D^b(\mathbf{A})$.*

Proof. The subcategory \mathbf{E} is closed under extensions, hence \mathbf{E} is a full exact subcategory of \mathbf{A} (see [21, Sect. 4]). Condition (E1) follows from the fact that, if f is an admissible epimorphism in \mathbf{E} , then $\ker f \in \mathbf{E}$. As $\mathcal{O}_{iZ} \in \mathbf{Perf}(X)$ for all $i > 0$, (E2) holds true taking $\{P_i\}_{i \in I} = \mathbf{Amp}(Z, X, H)$ defined in (2.3) (with H an ample divisor on X).

Obviously $D^b(\mathbf{E})$ is a full subcategory of $\mathbf{Perf}_Z(X)$. To show that they are actually equal, one has to apply an induction argument similar to the one in the first part of the proof of Proposition 3.7. To give a hint, let \mathbf{D}_n be the (strictly) full subcategory of $\mathbf{Perf}_Z(X)$ with objects the complexes \mathcal{A} with the following property: there exists $a \in \mathbb{Z}$ such that $H^p(\mathcal{A}) = 0$ for $p < a$ or $p > a + n$. Given $\mathcal{A} \in \mathbf{Perf}_Z(X)$, there exists $n \geq 0$ such that $\mathcal{A} \in \mathbf{D}_n$, and one can prove that $\mathcal{A} \in D^b(\mathbf{E})$ by induction on n . Indeed, if $n = 0$, there is nothing to prove. Otherwise we can assume without loss of generality that $H^p(\mathcal{A}) = 0$ for $p < -n$ or $p > 0$. Then \mathcal{A} sits in a distinguished triangle

$$\mathcal{C}[-1] \rightarrow P_i^{\oplus k} \rightarrow \mathcal{A} \rightarrow \mathcal{C},$$

where $P_i \in \mathbf{Amp}(Z, X, H)$, $k \in \mathbb{N}$ and $\mathcal{C} \in \mathbf{D}_{n-1}$.

As for (E3), one can prove more generally that for every $\mathcal{A} \in D^b(\mathbf{E}) = \mathbf{Perf}_Z(X)$ there exists an integer $N(\mathcal{A})$ such that $\mathrm{Hom}_{D^b(\mathbf{A})}(\mathcal{A}, \mathcal{B}[i]) = 0$, for every $i > N(\mathcal{A})$ and every $\mathcal{B} \in \mathbf{A}$. Indeed, this follows from the isomorphism

$$\mathrm{Hom}_{D^b(\mathbf{A})}(\mathcal{A}, \mathcal{B}[i]) \cong \mathrm{Hom}_{D^b(X)}(\mathcal{O}_X, \mathcal{A}^\vee \otimes \mathcal{B}[i]),$$

which holds because \mathcal{A} is perfect. More precisely, the cohomologies of $\mathcal{A}^\vee \otimes \mathcal{B}$ being bounded with bound depending only on \mathcal{A} , the vanishing of $\mathrm{Hom}_{D^b(X)}(\mathcal{O}_X, \mathcal{A}^\vee \otimes \mathcal{B}[i])$, for $i > N(\mathcal{A})$, can be deduced by induction on the cohomologies of $\mathcal{A}^\vee \otimes \mathcal{B}$ using the Grothendieck vanishing theorem (see, for example, [14, III, Thm. 2.7]). \square

Remark 3.11. In view of Proposition 2.12, it is easy to see that, if X, Z, \mathbf{E} and \mathbf{A} are as above, then condition $(*)$ in the introduction implies (Δ) . Indeed, in this case (1) in $(*)$ and (Δ) coincide. As for (2), consider $C \in D^b(\mathbf{E})$ with trivial cohomologies in positive degrees. Then, by Proposition 2.12, there is an integer N_1 such that for all $i < N_1$ and $j \ll i$ part (1) of Definition 2.9 holds true for $H^0(C)$ and $P_k := \mathcal{O}_{|i|Z}(jH) \in \mathbf{Amp}(Z, X, H)$, where $k = (i, j)$. By (2) in $(*)$, we can take another integer N_2 such that, for $i' < N_2$ and $j' \ll i'$,

$$\mathrm{Hom}(F(C), F(P_{k'})) = 0,$$

where $k' = (i', j')$. Considering $\min\{N_1, N_2\}$, this shows that (2) of (Δ) holds as well. Therefore, in the proof of Theorems 1.1 and 1.2 we can freely use the results in Section 3.3.

It may be useful to keep in mind some examples of exact functors satisfying $(*)$.

Example 3.12. In this example we assume that X_1 is a quasi-projective scheme with a projective subscheme Z_1 such that $\mathcal{O}_{iZ_1} \in \mathbf{Perf}(X_1)$, for all $i > 0$, and $T_0(\mathcal{O}_{Z_1}) = 0$.

(i) Using (2) in Definition 2.9 (which holds thanks to Proposition 2.12), it is very easy to verify that full functors $F: \mathbf{Perf}_{Z_1}(X_1) \rightarrow \mathbf{Perf}_{Z_2}(X_2)$ satisfy $(*)$ for any scheme X_2 containing a closed subscheme Z_2 .

(ii) For the same reason, a trivial example of a functor with the property $(*)$ but which is not full is $\text{id} \oplus \text{id}: \mathbf{Perf}_{Z_1}(X_1) \rightarrow \mathbf{Perf}_{Z_1}(X_1)$.

(iii) Following the same argument as in [11, Sect. 4], in the supported setting one may take exact functors $D_{Z_1}^b(X_1) \rightarrow D_{Z_2}^b(X_2)$ induced by exact full functors $\mathbf{Coh}_{Z_1}(X_1) \rightarrow \mathbf{Coh}_{Z_2}(X_2)$, where X_1 and X_2 are smooth quasi-projective varieties. These functors obviously satisfy $(*)$.

We conclude this section with the following easy result making clear that in the smooth case without support conditions, (\diamond) is equivalent to $(*)$ in the introduction.

Proposition 3.13. *Let X_1 be a smooth projective scheme such that $\dim(X_1) > 0$ and let X_2 be a scheme containing a closed subscheme Z_2 . Then an exact functor $F: D^b(X_1) \rightarrow D_{Z_2}^b(X_2)$ satisfies $(*)$ if and only if it satisfies (\diamond) .*

Proof. Clearly it is enough to show that (\diamond) implies (2) in $(*)$. Since $Z_1 = X_1$, $\mathcal{O}_{|i|Z_1}(jH_1) = \mathcal{O}_{X_1}(jH_1)$, for all $i, j \in \mathbb{Z}$, and when j varies they give rise to an almost ample set. Hence it is enough to show that for any $\mathcal{A} \in D^b(X_1)$ with trivial cohomologies in positive degrees, there is $N \in \mathbb{Z}$ such that $\text{Hom}(F(\mathcal{A}), F(\mathcal{O}_{X_1}(iH_1))) = 0$ for any $i < N$. Observing that F has a left adjoint F^* by [5] (see also [11, Rmk. 2.1]), the same argument as in the proof of Lemma 3.9 ensures that $F^* \circ F(\mathcal{A})$ has trivial cohomologies in positive degrees.

Assume, without loss of generality that $H^0(F^* \circ F(\mathcal{A}))$ is the last non-trivial cohomology. Then we have a distinguished triangle

$$\mathcal{A}' \rightarrow F^* \circ F(\mathcal{A}) \rightarrow H^0(F^* \circ F(\mathcal{A})) \rightarrow \mathcal{A}'[1]$$

where \mathcal{A}' has non-trivial cohomologies only in negative degrees. Hence $\text{Hom}(\mathcal{A}', \mathcal{B}) = 0$, for all $\mathcal{B} \in \mathbf{Coh}(X_1)$.

By the last statement in Proposition 2.12 (and (2) of Definition 2.9), there is an integer N such that, for all $i < N$ we have $\text{Hom}(H^0(F^* \circ F(\mathcal{A})), \mathcal{O}_{X_1}(iN_1)) = 0$. By the above remark and adjunction, we then have

$$0 = \text{Hom}(F^* \circ F(\mathcal{A}), \mathcal{O}_{X_1}(iN_1)) \cong \text{Hom}(F(\mathcal{A}), F(\mathcal{O}_{X_1}(iN_1))),$$

for all $i < N$. This is precisely (2) in $(*)$. □

Example 3.14. In view of Proposition 3.13 and of [11, Prop. 5.1], a non-trivial class of exact functors satisfying $(*)$ is provided by the functors $D^b(X_1) \rightarrow D^b(X_2)$ induced by exact functors $\mathbf{Coh}(X_1) \rightarrow \mathbf{Coh}(X_2)$. Here we assume that X_1 and X_2 are smooth projective varieties and that $\dim(X_1) > 0$.

As a consequence of Proposition 3.13, Theorem 1.1 generalizes the main result of [11] when the twists from the Brauer groups are trivial.

4. ENHANCEMENTS AND EXISTENCE OF FOURIER–MUKAI KERNELS

In this section we show how to construct Fourier–Mukai kernels for functors satisfying the condition $(*)$ defined in the introduction. This extends several results already present in the literature. Moreover we show that, in the supported setting, the Fourier–Mukai kernels have to be quasi-coherent rather than coherent. We need also to recall some basic facts about dg categories.

As an application of this machinery and of the results in the previous sections, we get the proof of Theorem 1.2. Here and for the rest of the paper, we fix a universe such that all dg categories are small dg categories with respect to this universe (see [28, Appendix A]).

4.1. Dg categories. In this section we give a quick introduction to some basic definitions and results about dg categories and dg functors. For a survey on the subject, the reader can have a look at [22].

Recall that a *dg category* is a \mathbb{k} -linear category \mathbf{A} such that, for all $A, B \in \text{Ob}(\mathbf{A})$, the morphism spaces $\text{Hom}(A, B)$ are \mathbb{Z} -graded \mathbb{k} -modules with a differential $d: \text{Hom}(A, B) \rightarrow \text{Hom}(A, B)$ of degree 1 and the composition maps are morphisms of complexes. Notice that the identity of each object is a closed morphism of degree 0.

Example 4.1. (i) Any \mathbb{k} -linear category has a (trivial) structure of dg category, with morphism spaces concentrated in degree 0.

(ii) For a dg category \mathbf{A} , one defines the opposite dg category \mathbf{A}° with $\text{Ob}(\mathbf{A}^\circ) = \text{Ob}(\mathbf{A})$ while $\text{Hom}_{\mathbf{A}^\circ}(A, B) := \text{Hom}_{\mathbf{A}}(B, A)$.

(iii) Following [12], given a dg category \mathbf{A} and a full dg subcategory \mathbf{B} of \mathbf{A} , one can form the quotient \mathbf{A}/\mathbf{B} which is again a dg category.

(iv) Given an abelian category \mathbf{A} , one can consider the dg category $\text{C}^{\text{dg}}(\mathbf{A})$ of complexes of objects in \mathbf{A} , its full dg subcategory $\text{Ac}^{\text{dg}}(\mathbf{A})$ of acyclic complexes and the dg quotient $\text{D}^{\text{dg}}(\mathbf{A}) := \text{C}^{\text{dg}}(\mathbf{A})/\text{Ac}^{\text{dg}}(\mathbf{A})$. When $\mathbf{A} = \mathbf{Qcoh}_Z(X)$, for X a scheme containing a closed subscheme Z , we denote the dg categories $\text{C}^{\text{dg}}(\mathbf{A})$, $\text{Ac}^{\text{dg}}(\mathbf{A})$ and $\text{D}^{\text{dg}}(\mathbf{A})$ respectively by $\text{C}_Z^{\text{dg}}(X)$, $\text{Ac}_Z^{\text{dg}}(X)$ and $\text{D}_Z^{\text{dg}}(X)$.

Given a dg category \mathbf{A} we denote by $H^0(\mathbf{A})$ its *homotopy category*. The objects of $H^0(\mathbf{A})$ are the same as those of \mathbf{A} while the morphisms from A to B are obtained by taking the 0-th cohomology $H^0(\text{Hom}_{\mathbf{A}}(A, B))$ of the complex $\text{Hom}_{\mathbf{A}}(A, B)$. If \mathbf{A} is pretriangulated (see [22] for the definition), then $H^0(\mathbf{A})$ has a natural structure of triangulated category.

Example 4.2. Given an abelian category \mathbf{A} , the dg categories $\text{C}^{\text{dg}}(\mathbf{A})$, $\text{Ac}^{\text{dg}}(\mathbf{A})$ and $\text{D}^{\text{dg}}(\mathbf{A})$ are pretriangulated and, as it is explained for example in [22, Sect. 4.4], there is an exact equivalence between the derived category $\text{D}(\mathbf{A})$ and the homotopy category $H^0(\text{D}^{\text{dg}}(\mathbf{A}))$. In particular, when X is a scheme containing a closed subscheme Z , the dg category $\text{D}_Z^{\text{dg}}(X)$ is pretriangulated and $H^0(\text{D}_Z^{\text{dg}}(X)) \cong \text{D}_Z(\mathbf{Qcoh}(X))$ (here one uses that $\text{D}_Z(\mathbf{Qcoh}(X))$ is naturally equivalent to $\text{D}(\mathbf{Qcoh}_Z(X))$ by Proposition 2.7).

A *dg functor* $F: \mathbf{A} \rightarrow \mathbf{B}$ between two dg categories is the datum of a map $\text{Ob}(\mathbf{A}) \rightarrow \text{Ob}(\mathbf{B})$ and of morphisms of complexes of \mathbb{k} -modules $\text{Hom}_{\mathbf{A}}(A, B) \rightarrow \text{Hom}_{\mathbf{B}}(F(A), F(B))$, for $A, B \in \text{Ob}(\mathbf{A})$, which are compatible with the compositions and the units.

A dg functor $F: \mathbf{A} \rightarrow \mathbf{B}$ induces a functor $H^0(F): H^0(\mathbf{A}) \rightarrow H^0(\mathbf{B})$, which is exact (between triangulated categories) if \mathbf{A} and \mathbf{B} are pretriangulated.

A dg functor $F: \mathbf{A} \rightarrow \mathbf{B}$ is a *quasi-equivalence*, if the maps $\text{Hom}(A, B) \rightarrow \text{Hom}(F(A), F(B))$ are quasi-isomorphisms, for every $A, B \in \mathbf{A}$, and $H^0(F)$ is an equivalence. One can consider the localization \mathbf{Hqe} of the category of dg categories over \mathbb{k} with respect to quasi-equivalences ([32]).

Given a dg functor F , we will denote with the same symbol its image in \mathbf{Hqe} . In particular, if F is a quasi-equivalence, we denote by F^{-1} the morphism in \mathbf{Hqe} which is the inverse of F .

For a small dg category \mathbf{A} , one can consider the pretriangulated dg category $\mathcal{M}od\text{-}\mathbf{A}$ of *right dg \mathbf{A} -modules*. A right dg \mathbf{A} -module is a dg functor $M: \mathbf{A}^\circ \rightarrow \mathcal{M}od\text{-}\mathbb{k}$, where $\mathcal{M}od\text{-}\mathbb{k}$ is the dg category of dg \mathbb{k} -modules. The full dg subcategory of acyclic right dg modules is denoted by $\mathcal{A}c(\mathbf{A})$, and $H^0(\mathcal{A}c(\mathbf{A}))$ is a full triangulated subcategory of the homotopy category $H^0(\mathcal{M}od\text{-}\mathbf{A})$. Hence the *derived category* of the dg category \mathbf{A} is the Verdier quotient

$$\mathcal{D}(\mathbf{A}) := H^0(\mathcal{M}od\text{-}\mathbf{A})/H^0(\mathcal{A}c(\mathbf{A})).$$

A right dg \mathbf{A} -module is *representable* if it is contained in the image of the Yoneda dg functor

$$Y: \mathbf{A} \rightarrow \mathcal{M}od\text{-}\mathbf{A} \quad A \mapsto \text{Hom}_{\mathbf{A}}(-, A) =: Y^A.$$

A right dg \mathbf{A} -module is *free* if it is isomorphic to a direct sum of dg modules of the form $Y^A[m]$, where $A \in \mathbf{A}$ and $m \in \mathbb{Z}$. A right dg \mathbf{A} -module M is *semi-free* if it has a filtration

$$0 = M_0 \subseteq M_1 \subseteq \dots = M$$

such that M_i/M_{i-1} is free, for all i . We denote by $\text{SF}(\mathbf{A})$ the full dg subcategory of semi-free dg modules, while $\text{SF}_{\text{fg}}(\mathbf{A}) \subseteq \text{SF}(\mathbf{A})$ is the full dg subcategory of *finitely generated semi-free* dg modules. Namely, there is n such that $M_n = M$ and each M_i/M_{i-1} is a finite direct sum of dg modules of the form $Y^A[m]$. The dg modules which are homotopy equivalent to direct summands of finitely generated semi-free dg modules are called *perfect* and they form a full dg subcategory $\text{Perf}^{\text{dg}}(\mathbf{A})$.

Following [22, 32], given two dg categories \mathbf{A} and \mathbf{B} , we denote by $\text{rep}(\mathbf{A}, \mathbf{B})$ the full subcategory of the derived category $\mathcal{D}(\mathbf{A}^\circ \otimes_{\mathbb{k}} \mathbf{B})$ of \mathbf{A} - \mathbf{B} -bimodules C such that the functor $(-)\otimes_{\mathbf{A}} C: \mathcal{D}(\mathbf{A}) \rightarrow \mathcal{D}(\mathbf{B})$ sends the representable \mathbf{A} -modules to objects which are isomorphic to representable \mathbf{B} -modules. An object in $\text{rep}(\mathbf{A}, \mathbf{B})$ is called a *quasi-functor*. By [32], morphisms in \mathbf{Hqe} are in natural bijection with isomorphism classes of quasi-functors. Thus, with a slight abuse of notation, we sometimes call quasi-functor a morphism in \mathbf{Hqe} . Notice that a quasi-functor $M \in \text{rep}(\mathbf{A}, \mathbf{B})$ induces a functor $H^0(M): H^0(\mathbf{A}) \rightarrow H^0(\mathbf{B})$, well defined up to isomorphism.

For $F: \mathbf{A} \rightarrow \mathbf{B}$ a dg functor, there exist dg functors

$$F^*: \mathcal{M}od\text{-}\mathbf{A} \rightarrow \mathcal{M}od\text{-}\mathbf{B} \quad F_*: \mathcal{M}od\text{-}\mathbf{B} \rightarrow \mathcal{M}od\text{-}\mathbf{A}$$

also denoted, respectively, by Ind_F and Res_F . While F_* is simply induced by composition with F , the reader can have a look at [12, Sect. 14] for the definition and properties of F^* . In particular, F^* is left adjoint to F_* and commutes with the Yoneda embeddings, up to dg isomorphism. Moreover, F^* preserves semi-free dg modules and $F^*: \text{SF}(\mathbf{A}) \rightarrow \text{SF}(\mathbf{B})$ is a quasi-equivalence if $F: \mathbf{A} \rightarrow \mathbf{B}$ is such.

Given two pretriangulated dg categories \mathbf{A} and \mathbf{B} and an exact functor $F: H^0(\mathbf{A}) \rightarrow H^0(\mathbf{B})$, a *dg lift* of F is a quasi-functor $G \in \text{rep}(\mathbf{A}, \mathbf{B})$ such that $H^0(G) \cong F$.

An *enhancement* of a triangulated category \mathbf{T} is a pair (\mathbf{A}, α) , where \mathbf{A} is a pretriangulated dg category and $\alpha: H^0(\mathbf{A}) \rightarrow \mathbf{T}$ is an exact equivalence. The enhancement (\mathbf{A}, α) of \mathbf{T} is *unique* if for any enhancement (\mathbf{B}, β) of \mathbf{T} there exists a quasi-functor $\gamma: \mathbf{A} \rightarrow \mathbf{B}$ such that $H^0(\gamma): H^0(\mathbf{A}) \rightarrow H^0(\mathbf{B})$ is an exact equivalence. We say that the enhancement is *strongly unique* if moreover γ can

be chosen so that $\alpha \cong \beta \circ H^0(\gamma)$. Often, by abuse of notation, we will say that a pretriangulated dg category \mathbf{A} is an enhancement of a triangulated category \mathbf{T} when there is an exact equivalence $H^0(\mathbf{A}) \cong \mathbf{T}$.

Example 4.3. If \mathbf{A} is a dg category, $\mathrm{SF}(\mathbf{A})$ and $\mathrm{Perf}^{\mathrm{dg}}(\mathbf{A})$ are enhancements, respectively, of $\mathcal{D}(\mathbf{A})$ and $\mathcal{D}(\mathbf{A})^c$.

Example 4.4. Let \mathbf{A} be a pretriangulated dg category and \mathbf{B} a full pretriangulated dg subcategory of \mathbf{A} . By [12], there exists a natural exact equivalence between the Verdier quotient $H^0(\mathbf{A})/H^0(\mathbf{B})$ and $H^0(\mathbf{A}/\mathbf{B})$. Hence \mathbf{A}/\mathbf{B} is an enhancement of $H^0(\mathbf{A})/H^0(\mathbf{B})$.

Example 4.5. By Example 4.2, $\mathrm{D}^{\mathrm{dg}}(\mathbf{A})$ (for \mathbf{A} an abelian category) is an enhancement of $\mathrm{D}(\mathbf{A})$. Moreover, if X is a scheme containing a closed subscheme Z , $\mathrm{D}_Z^{\mathrm{dg}}(X)$ is an enhancement of $\mathrm{D}_Z(\mathbf{Qcoh}(X))$. Let $\mathrm{Perf}_Z^{\mathrm{dg}}(X)$ be the full dg subcategory of $\mathrm{D}_Z^{\mathrm{dg}}(X)$ consisting of compact objects in $H^0(\mathrm{D}_Z^{\mathrm{dg}}(X))$. Notice that $\mathrm{Perf}_Z^{\mathrm{dg}}(X)$ is an enhancement of $\mathbf{Perf}_Z(X)$ and, as we mentioned above, we will identify $H^0(\mathrm{Perf}_Z^{\mathrm{dg}}(X))$ with $\mathbf{Perf}_Z(X)$, to make the notation simpler.

4.2. Enhancements and the proof of Theorem 1.2. Let X be a quasi-projective scheme containing a projective subscheme Z and let H be an ample divisor on X . Assume that $\mathcal{O}_{iZ} \in \mathbf{Perf}(X)$ for all $i > 0$ (hence the full subcategory $\mathbf{Amp}(Z, X, H)$ defined in (2.3) is contained in $\mathbf{Perf}_Z(X)$). Consider the dg category $\mathbf{A} := \mathbf{Coh}_Z(X) \cap \mathbf{Perf}_Z(X)$ concentrated in degree zero and notice that, due to Remark 2.10 and Proposition 2.12, the objects in \mathbf{A} satisfy (1) in Definition 2.9. By abuse of notation, we will write \mathbf{Y} for the functor $\mathbf{A} \rightarrow \mathcal{D}(\mathbf{A})$ which is the composition of $H^0(\mathbf{Y}): \mathbf{A} = H^0(\mathbf{A}) \rightarrow H^0(\mathrm{Mod}\text{-}\mathbf{A})$ and of the quotient functor $H^0(\mathrm{Mod}\text{-}\mathbf{A}) \rightarrow \mathcal{D}(\mathbf{A})$. As a matter of notation, if \mathbf{T} is a triangulated category with arbitrary direct sums and \mathbf{L} is a localizing subcategory of \mathbf{T} , we denote by $\pi: \mathbf{T} \rightarrow \mathbf{T}/\mathbf{L}$ the quotient functor. Recall that a full triangulated subcategory \mathbf{S} of a triangulated category \mathbf{T} is *localizing* if it is closed under arbitrary direct sums.

Lemma 4.6. *There exists an exact equivalence $\varphi: \mathrm{D}_Z(\mathbf{Qcoh}(X)) \rightarrow \mathcal{D}(\mathbf{A})/\mathbf{L}$, for some localizing subcategory $\mathbf{L} \subseteq \mathcal{D}(\mathbf{A})$, such that we have an isomorphism of functors $\pi \circ \mathbf{Y} \cong \varphi|_{\mathbf{A}}$. Moreover $\mathrm{D}_Z(\mathbf{Qcoh}(X))$ has a unique enhancement.*

Proof. By Proposition 2.12, the category $\mathbf{Amp}(Z, X, H) \subseteq \mathbf{A}$ is a set of compact generators for the Grothendieck category $\mathbf{Qcoh}_Z(X)$. Then take the abelian category $\mathrm{Mod}\text{-}\mathbf{A}$ of modules over \mathbf{A} , i.e. \mathbb{k} -linear contravariant functors from \mathbf{A} to the category of \mathbb{k} -modules. As it is explained in [8] there is a Serre subcategory \mathbf{N} of $\mathrm{Mod}\text{-}\mathbf{A}$ such that $\mathbf{Qcoh}_Z(X) \cong \mathrm{Mod}\text{-}\mathbf{A}/\mathbf{N}$. By [28, Lemma 7.2], we then have an equivalence

$$(4.1) \quad \mathrm{D}(\mathbf{Qcoh}_Z(X)) \cong \mathrm{D}(\mathrm{Mod}\text{-}\mathbf{A})/\mathrm{D}_{\mathbf{N}}(\mathrm{Mod}\text{-}\mathbf{A}),$$

where $\mathrm{D}_{\mathbf{N}}(\mathrm{Mod}\text{-}\mathbf{A})$ is the full subcategory of $\mathrm{D}(\mathrm{Mod}\text{-}\mathbf{A})$ consisting of complexes with cohomologies in \mathbf{N} . As observed at the beginning of Section 7 in [28], there exists a natural equivalence $\psi: \mathrm{D}(\mathrm{Mod}\text{-}\mathbf{A}) \rightarrow \mathcal{D}(\mathbf{A})$. Hence we set \mathbf{L} to be the full subcategory of $\mathcal{D}(\mathbf{A})$ corresponding to $\mathrm{D}_{\mathbf{N}}(\mathrm{Mod}\text{-}\mathbf{A})$ under ψ .

We define φ to be the composition of (4.1) with the equivalences $\mathrm{D}(\mathrm{Mod}\text{-}\mathbf{A})/\mathrm{D}_{\mathbf{N}}(\mathrm{Mod}\text{-}\mathbf{A}) \cong \mathcal{D}(\mathbf{A})/\mathbf{L}$ (induced by ψ) and $\mathrm{D}_Z(\mathbf{Qcoh}(X)) \cong \mathrm{D}(\mathbf{Qcoh}_Z(X))$ (see Proposition 2.7). The fact that there is an isomorphism of functors $\pi \circ \mathbf{Y} \cong \varphi|_{\mathbf{A}}$ is observed in [8], where the above construction

is analyzed further. Notice that the Yoneda embedding $\mathbf{Y}: \mathbf{A} \rightarrow \mathbf{D}(\mathbf{A})$ coincides with the classical Yoneda embedding $\mathbf{A} \rightarrow \mathbf{Mod}\text{-}\mathbf{A}$, composed with the natural inclusion $\mathbf{Mod}\text{-}\mathbf{A} \hookrightarrow \mathbf{D}(\mathbf{Mod}\text{-}\mathbf{A})$ and with ψ .

The second part of the statement is a straightforward consequence of [28, Thm. 7.5]. \square

As a consequence, we have an equivalence

$$(4.2) \quad \alpha: \mathbf{Perf}_Z(X) \rightarrow (\mathcal{D}(\mathbf{A})/\mathbf{L})^c$$

induced by φ (i.e. we set $\alpha := \varphi|_{\mathbf{Perf}_Z(X)}$). This is because φ , being an equivalence, sends compact objects to compact objects. By Lemma 4.6, we have an isomorphism of functors $\alpha^{-1} \circ \pi \circ \mathbf{Y} \cong \text{id}_{\mathbf{A}}$ (where α^{-1} denotes a quasi-inverse of α).

Let \mathbf{L}' be the lift of the localizing subcategory \mathbf{L} to $\mathcal{M}od\text{-}\mathbf{A}$. Let \mathbf{D} be the full dg subcategory of $\mathbf{SF}(\mathbf{A})/(\mathbf{SF}(\mathbf{A}) \cap \mathbf{L}')$ (which, by Examples 4.3 and 4.4, is an enhancement of $\mathcal{D}(\mathbf{A})/\mathbf{L}$) consisting of the compact objects in $H^0(\mathbf{SF}(\mathbf{A})/(\mathbf{SF}(\mathbf{A}) \cap \mathbf{L}'))$. Obviously, \mathbf{D} is an enhancement of $(\mathcal{D}(\mathbf{A})/\mathbf{L})^c$ in a natural way. In view of this, we will identify $H^0(\mathbf{D})$ with $(\mathcal{D}(\mathbf{A})/\mathbf{L})^c$.

Lemma 4.7. *If (\mathbf{B}, β) is an enhancement of $\mathbf{Perf}_Z(X)$, there exists a quasi-functor $\delta: \mathbf{D} \rightarrow \mathbf{B}$ such that $H^0(\delta)$ is an exact equivalence and there is an isomorphism of functors $\mathbf{A} \rightarrow H^0(\mathbf{B})$*

$$(4.3) \quad H^0(\delta) \circ \pi \circ \mathbf{Y} \xrightarrow{\sim} (\alpha \circ \beta)^{-1} \circ \pi \circ \mathbf{Y}.$$

Proof. Due to [8], the category \mathbf{L} is compactly generated in $\mathcal{D}(\mathbf{A}) \cong \mathbf{D}(\mathbf{Mod}\text{-}\mathbf{A})$ and $\mathbf{L}^c = \mathbf{L} \cap \mathcal{D}(\mathbf{A})^c$. Hence we can apply [28, Thm. 6.4] to the exact equivalence $(\alpha \circ \beta)^{-1}$, providing the quasi-functor $\delta: \mathbf{D} \rightarrow \mathbf{B}$ in the statement. Indeed, $H^0(\delta)$ is fully faithful (by (1) in [28, Thm. 6.4]), satisfies (4.3) (by (2) in [28, Thm. 6.4]) and is essentially surjective (by (3) in [28, Thm. 6.4] and the fact that $(\alpha \circ \beta)^{-1}$ is an equivalence). \square

Now we want to prove Theorem 1.2 and so we assume further that $T_0(\mathcal{O}_Z) = 0$. Let us reproduce the statement here for the convenience of the reader.

Theorem 4.8. *Let X be a quasi-projective scheme containing a projective subscheme Z such that $\mathcal{O}_{iZ} \in \mathbf{Perf}(X)$, for all $i > 0$, and $T_0(\mathcal{O}_Z) = 0$. Then $\mathbf{Perf}_Z(X)$ has a strongly unique enhancement.*

Proof. Let (\mathbf{B}, β) be an enhancement of $\mathbf{Perf}_Z(X)$. By Lemma 4.7, there exists a quasi-functor $\delta: \mathbf{D} \rightarrow \mathbf{B}$ such that $H^0(\delta)$ is an exact equivalence satisfying (4.3).

By Proposition 3.10, there is a natural exact equivalence $\varepsilon: \mathbf{D}^b(\mathbf{E}) \rightarrow \mathbf{Perf}_Z(X)$, for the exact category $\mathbf{E} = \mathbf{Perf}_Z(X) \cap \mathbf{Coh}_Z(X)$. Setting $\varpi := \alpha \circ \varepsilon$, we have $\varpi^{-1} \circ \pi \circ \mathbf{Y} \cong \text{id}_{\mathbf{A}}$.

Put $\mathbf{F}_1 := H^0(\delta) \circ \varpi$ and $\mathbf{F}_2 := (\alpha \circ \beta)^{-1} \circ \varpi$. Then (4.3) reads as

$$(4.4) \quad \mathbf{F}_1|_{\mathbf{A}} \xrightarrow{\sim} \mathbf{F}_2|_{\mathbf{A}}.$$

By Corollary 3.8, it extends to a unique isomorphism $\mathbf{F}_1 \cong \mathbf{F}_2$. Notice that this is the point where we use that $T_0(\mathcal{O}_Z) = 0$ as, under this assumption, \mathbf{A} is an almost ample set (by Proposition 2.12 and Remark 2.10) and, using this, every full functor certainly satisfies (Δ) (see Remark 3.11 and Example 3.12). Therefore there exists an isomorphism between $H^0(\delta)$ and $(\alpha \circ \beta)^{-1}$.

This proves that the enhancement of $\mathbf{Perf}_Z(X)$ is strongly unique. Indeed, suppose that (\mathbf{B}_1, β_1) and (\mathbf{B}_2, β_2) are enhancements of $(\mathcal{D}(\mathbf{A})/\mathbf{L})^c$. By the above discussion, there are quasi-equivalences

$\delta_i: \mathbf{D} \rightarrow \mathbf{B}_i$ and unique isomorphisms $H^0(\delta_i) \cong \beta_i^{-1} \circ \alpha^{-1}$. To conclude, if we set $\tilde{\delta} := \delta_2 \circ \delta_1^{-1}: \mathbf{B}_1 \rightarrow \mathbf{B}_2$, we have $\beta_2 \circ H^0(\tilde{\delta}) \cong \beta_1$. \square

In view of Example 2.11, it is straightforward to deduce the following special instance of Theorem 1.2.

Corollary 4.9. *Let X be a quasi-projective scheme containing a projective subscheme Z such that $T_0(\mathcal{O}_Z) = 0$ and either X is smooth or $X = Z$. Then $\mathbf{Perf}_Z(X)$ has a strongly unique enhancement.*

If $X = Z$, then this is nothing but one of the main results in [28] (see Theorem 9.9 there).

4.3. The Fourier–Mukai kernels. The aim of this subsection is to prove the following result, which is the first part of Theorem 1.1.

Proposition 4.10. *Let X_1 be a quasi-projective scheme containing a projective subscheme Z_1 such that $\mathcal{O}_{iZ_1} \in \mathbf{Perf}(X_1)$, for all $i > 0$. Assume that X_2 is a scheme containing a closed subscheme Z_2 . Then, for any exact functor $F: \mathbf{Perf}_{Z_1}(X_1) \rightarrow \mathbf{Perf}_{Z_2}(X_2)$ satisfying $(*)$ there exist $\mathcal{E} \in \mathbf{D}_{Z_1 \times Z_2}(\mathbf{Qcoh}(X_1 \times X_2))$ and an isomorphism of exact functors $F \cong \Phi_{\mathcal{E}}^s$.*

To construct the Fourier–Mukai kernel realizing F as a Fourier–Mukai functor we will make use of some ideas from Sections 4, 6 and 9 of [28]. Clearly there is no space for reproducing the arguments in [28] in full details. Thus, for the convenience of the reader, we outline here the main steps in the proof of Proposition 4.10.

The first passage consists in the construction of a quasi-functor F_1^{dg} which will turn out to be a dg lift of F . This is done in Section 4.3.1 following [28, Sect. 4]. Then we will show, in Section 4.3.2, the existence of the above mentioned isomorphism. The first main ingredient at this point is that the dg enhancement of the category of perfect complexes with support conditions is constructed out of a set satisfying (1) of Definition 2.9. The second ingredient consists in showing that F and $F_1 := H^0(F_1^{\mathrm{dg}})$ are isomorphic on such a set. For this we need to apply some tricks from [28, Sect. 6]. Hence, using that the functor F satisfies $(*)$, the results in Section 3 apply, proving that $F \cong F_1$. Finally, we prove in Section 4.3.3 that F_1 is of Fourier–Mukai type. This essentially follows, as in [28, Sect. 9], from an application of [32].

We will discuss the relevant points where one can avoid the fully faithful assumption on F which is present in [28]. Anyway, it is important to observe that the steps in the following proof where such an assumption may be relevant are where we make use of Section 6 in [28]. On the other hand, only two results from that part of [28] are used here: Lemmas 6.1 and 6.2. While the first one is absolutely innocuous, one needs to observe that $(*)$ (together with [8]) is enough to apply [28, Lemma 6.2].

4.3.1. The dg lift. We keep the same notation as in Section 4.2, with X_1 and Z_1 playing the same roles as X and Z there. Set $F' := F \circ \alpha^{-1}$, where α is defined in (4.2). Since the essential image of $\pi \circ \mathbf{Y}$ is contained in $(\mathcal{D}(\mathbf{A})/\mathbf{L})^c$, we can define the full dg subcategory $\mathbf{B} := \{F'(\pi(\mathbf{Y}^A)) : A \in \mathbf{A}\}$ of $\mathbf{Perf}_{Z_2}^{\mathrm{dg}}(X_2)$ (see Example 4.5).

Let \mathbf{C} be the full dg subcategory of $\mathbf{Perf}_{Z_2}^{\mathrm{dg}}(X_2)$ such that $H^0(\mathbf{C})$ is classically generated by the objects in \mathbf{B} . By definition the functor $F': (\mathcal{D}(\mathbf{A})/\mathbf{L})^c \rightarrow \mathbf{Perf}_{Z_2}(X_2)$ factors through $H^0(\mathbf{C})$

giving rise to the functor

$$\mathbf{A} \xrightarrow{\pi \circ \mathcal{Y}} (\mathcal{D}(\mathbf{A})/\mathbf{L})^c \xrightarrow{F'} H^0(\mathbf{C}) \hookrightarrow \mathbf{Perf}_{Z_2}(X_2)$$

which, in turn, factors through a functor $\mathbf{G}_1: \mathbf{A} \rightarrow H^0(\mathbf{B})$, which can be viewed as a dg functor in a trivial way.

Consider the dg category $\tau_{\leq 0}\mathbf{B}$ with the same objects as \mathbf{B} but such that $\mathrm{Hom}_{\tau_{\leq 0}\mathbf{B}}(\mathcal{E}, \mathcal{F}) = \tau_{\leq 0}\mathrm{Hom}_{\mathbf{B}}(\mathcal{E}, \mathcal{F})$ (here $\tau_{\leq 0}$ is the gentle truncation). Let $p: \tau_{\leq 0}\mathbf{B} \rightarrow H^0(\mathbf{B})$ and $l: \tau_{\leq 0}\mathbf{B} \rightarrow \mathbf{B}$ be the natural dg functors. Due to Lemma 4.6 and part (1) of (*), p is a quasi-equivalence. Thus, we get the quasi-functor

$$\mathbf{G}_2: \mathrm{SF}(\mathbf{A}) \rightarrow \mathrm{SF}(H^0(\mathbf{B})) \rightarrow \mathrm{SF}(\tau_{\leq 0}\mathbf{B}) \rightarrow \mathrm{SF}(\mathbf{B}),$$

where $\mathbf{G}_2 = l^* \circ (p^*)^{-1} \circ \mathbf{G}_1^*$ and $(p^*)^{-1}$ is the inverse in \mathbf{Hqe} of p^* .

Since $\mathbf{L} = H^0(\mathbf{L}')$ is compactly generated by [8], the argument in [28, Lemma 6.2] applies and the quasi-functor \mathbf{G}_2 factors through the dg quotient $\mathrm{SF}(\mathbf{A})/(\mathrm{SF}(\mathbf{A}) \cap \mathbf{L}')$. Hence we get a quasi-functor

$$\mathbf{G}_3: \mathrm{SF}(\mathbf{A})/(\mathrm{SF}(\mathbf{A}) \cap \mathbf{L}') \rightarrow \mathrm{SF}(\mathbf{B}).$$

It is important to stress here that, in the proof of [28, Lemma 6.2], the assumption that \mathbf{F} is fully faithful is not needed and (*) suffices to conclude. This is because the proof relies on [28, Prop. 3.4], where part (1) of (*) is enough.

The quasi-functor \mathbf{G}_3 restricts to a quasi-functor $\mathbf{G}_3: \mathbf{D} \rightarrow \mathrm{Perf}^{\mathrm{dg}}(\mathbf{B})$. Indeed, it is enough to observe that by definition $H^0(\mathbf{G}_3)(\pi(\mathcal{Y}^A)) \in H^0(\mathrm{Perf}^{\mathrm{dg}}(\mathbf{B}))$, for all $A \in \mathbf{A}$. Now, [28, Prop. 1.16] applies and gives a quasi-equivalence $\psi: \mathbf{C} \rightarrow \mathrm{Perf}^{\mathrm{dg}}(\mathbf{B})$. Take the quasi-functor

$$(4.5) \quad \mathbf{G}_4: \mathbf{D} \xrightarrow{\mathbf{G}_3} \mathrm{Perf}^{\mathrm{dg}}(\mathbf{B}) \xrightarrow{\psi^{-1}} \mathbf{C} \hookrightarrow \mathrm{Perf}_{Z_2}^{\mathrm{dg}}(X_2),$$

where, as usual, ψ^{-1} is the inverse in \mathbf{Hqe} of the quasi-equivalence ψ .

We apply Lemma 4.7 to the enhancement $\mathrm{Perf}_{Z_1}^{\mathrm{dg}}(X_1)$ of $\mathbf{Perf}_{Z_1}(X_1)$ getting a quasi-functor $\delta: \mathbf{D} \rightarrow \mathrm{Perf}_{Z_1}^{\mathrm{dg}}(X_1)$, whence

$$\mathbf{F}_1^{\mathrm{dg}} := \mathbf{G}_4 \circ \delta^{-1}: \mathrm{Perf}_{Z_1}^{\mathrm{dg}}(X_1) \longrightarrow \mathrm{Perf}_{Z_2}^{\mathrm{dg}}(X_2).$$

4.3.2. *The isomorphism.* Consider the exact functor

$$H^0(\mathbf{G}_4): (\mathcal{D}(\mathbf{A})/\mathbf{L})^c (= H^0(\mathbf{D})) \longrightarrow \mathbf{Perf}_{Z_2}(X_2).$$

As a consequence of [28, Lemma 6.1] (see also [28, Prop. 3.4]) and of the definition in (4.5), we get an isomorphism

$$\theta: F' \circ \pi \circ \mathcal{Y} \xrightarrow{\sim} H^0(\mathbf{G}_4) \circ \pi \circ \mathcal{Y},$$

as functors from \mathbf{A} to $\mathbf{Perf}_{Z_2}(X_2)$. Again, it is important to observe that we do not need to have \mathbf{F} (and thus F') fully faithful to apply [28, Lemma 6.1] being the isomorphism above a simple consequence of the definition of \mathbf{G}_4 .

From the isomorphisms (4.3) and $\alpha^{-1} \circ \pi \circ \mathcal{Y} \cong \mathrm{id}_{\mathbf{A}}$ (see Lemma 4.6), θ gives an isomorphism

$$\mathbf{F}|_{\mathbf{A}} \xrightarrow{\sim} \mathbf{F}_1|_{\mathbf{A}},$$

where $\mathbf{F}_1 := H^0(\mathbf{F}_1^{\mathrm{dg}})$. Applying Corollary 3.8 we get an isomorphism of exact functors

$$(4.6) \quad \mathbf{F} \xrightarrow{\sim} \mathbf{F}_1.$$

4.3.3. *The kernel.* By [28, Prop. 1.17] (which can be applied here in view of Proposition 2.1), we get a quasi-equivalence $\varphi_i: \mathbf{SF}(\mathbf{Perf}_{Z_i}^{\mathrm{dg}}(X_i)) \xrightarrow{\sim} \mathbf{D}_{Z_i}^{\mathrm{dg}}(X_i)$. Hence, composing the extension of $\mathbf{F}_1^{\mathrm{dg}}$ to semi-free modules with φ_i , we get a quasi-functor

$$\mathbf{F}_2^{\mathrm{dg}}: \mathbf{D}_{Z_1}^{\mathrm{dg}}(X_1) \xrightarrow{\varphi_1^{-1}} \mathbf{SF}(\mathbf{Perf}_{Z_1}^{\mathrm{dg}}(X_1)) \rightarrow \mathbf{SF}(\mathbf{Perf}_{Z_2}^{\mathrm{dg}}(X_2)) \xrightarrow{\varphi_2} \mathbf{D}_{Z_2}^{\mathrm{dg}}(X_2)$$

whose H^0 commutes with direct sums (because it has a right adjoint, according to [28, Sect. 1]). Observe that

$$(4.7) \quad \mathbf{F}_1 \cong H^0(\mathbf{F}_2^{\mathrm{dg}})|_{\mathbf{Perf}_{Z_1}(X_1)}.$$

The easier case $X_i = Z_i$, for $i = 1, 2$, generalizing [28, Cor. 9.13] (see, in particular, parts (2) and (3) there) can be treated already.

Corollary 4.11. *Let X_1 be a projective variety and let X_2 be a scheme. For any exact functor $\mathbf{F}: \mathbf{Perf}(X_1) \rightarrow \mathbf{Perf}(X_2)$ satisfying $(*)$, there exist $\mathcal{E} \in \mathbf{D}^b(X_1 \times X_2)$ and an isomorphism of exact functors $\mathbf{F} \cong \Phi_{\mathcal{E}}$.*

Proof. By [32, Thm. 8.9] there is $\mathcal{E} \in \mathbf{D}(\mathbf{Qcoh}(X_1 \times X_2))$ such that $\Phi_{\mathcal{E}} \cong H^0(\mathbf{F}_2^{\mathrm{dg}})$. As $\mathbf{F}_1 \cong H^0(\mathbf{F}_1^{\mathrm{dg}}) \cong H^0(\mathbf{F}_2^{\mathrm{dg}})|_{\mathbf{Perf}(X_1)}$, the isomorphism (4.6) gives $\mathbf{F} \cong \Phi_{\mathcal{E}}$. The fact that \mathcal{E} is bounded coherent is obtained by the same argument as in the proof of [28, Cor. 9.13], part (4). We do not explain this here as this is a special instance of Lemma 5.4. \square

Back to the general setting, we can observe the following.

Remark 4.12. Assume that X is a quasi-projective scheme containing a projective subscheme Z such that $\mathcal{O}_{iZ} \in \mathbf{Perf}(X)$, for all $i > 0$. The exact functors $\iota: \mathbf{D}_Z(\mathbf{Qcoh}(X)) \rightarrow \mathbf{D}(\mathbf{Qcoh}(X))$ and $\iota^!: \mathbf{D}(\mathbf{Qcoh}(X)) \rightarrow \mathbf{D}_Z(\mathbf{Qcoh}(X))$, defined in Section 2.1, have natural dg lifts (denoted with the same symbols) $\iota: \mathbf{D}_Z^{\mathrm{dg}}(X) \rightarrow \mathbf{D}^{\mathrm{dg}}(X)$ and $\iota^!: \mathbf{D}^{\mathrm{dg}}(X) \rightarrow \mathbf{D}_Z^{\mathrm{dg}}(X)$.

To show this one can use [25, Sect. 4]. More precisely, the presence of the right adjoint $\iota^!$ yields a semi-orthogonal decomposition of $\mathbf{D}(\mathbf{Qcoh}(X))$, where $\mathbf{D}_Z(\mathbf{Qcoh}(X))$ is a non-trivial piece. By [25, Prop. 4.10], such a decomposition can be written at the dg level as a gluing with one piece given by $\mathbf{D}_Z^{\mathrm{dg}}(X)$, up to quasi-equivalence. Hence [25, Lemma 4.4] combined with the uniqueness of the enhancements of $\mathbf{D}(\mathbf{Qcoh}(X))$ (see [28, Cor. 2.11]) and of $\mathbf{D}_Z(\mathbf{Qcoh}(X))$ (see Lemma 4.6) provides the dg lifts for the functors ι and $\iota^!$. By the construction, it is clear that the dg lifts of ι and $\iota^!$ are not, in general, dg functors but just quasi-functors.

In light of the above remark, consider the quasi-functor $\mathbf{F}_3^{\mathrm{dg}}$ making the following diagram commutative

$$(4.8) \quad \begin{array}{ccc} \mathbf{D}^{\mathrm{dg}}(X_1) & \xrightarrow{\mathbf{F}_3^{\mathrm{dg}}} & \mathbf{D}^{\mathrm{dg}}(X_2) \\ \iota_1^! \downarrow & & \uparrow \iota_2 \\ \mathbf{D}_{Z_1}^{\mathrm{dg}}(X_1) & \xrightarrow{\mathbf{F}_2^{\mathrm{dg}}} & \mathbf{D}_{Z_2}^{\mathrm{dg}}(X_2). \end{array}$$

Clearly $H^0(\mathbf{F}_3^{\mathrm{dg}})$ commutes with direct sums, as the same is true for $\iota_1^!$ (by Lemma 2.3), $H^0(\mathbf{F}_2^{\mathrm{dg}})$ and ι_2 . Notice also that, if $Z_1 = X_1$, then $\iota_1^! \cong \mathrm{id}$.

By [32, Thm. 8.9], there exist $\tilde{\mathcal{E}} \in D(\mathbf{Qcoh}(X_1 \times X_2))$ and an isomorphism of exact functors

$$\Phi_{\tilde{\mathcal{E}}} \cong H^0(\mathbf{F}_3^{\mathrm{dg}}): D(\mathbf{Qcoh}(X_1)) \xrightarrow{\iota_1^!} D_{Z_1}(\mathbf{Qcoh}(X_1)) \xrightarrow{H^0(\mathbf{F}_2^{\mathrm{dg}})} D_{Z_2}(\mathbf{Qcoh}(X_2)) \xrightarrow{\iota_2} D(\mathbf{Qcoh}(X_2)).$$

It follows that, setting $\mathcal{E} := \iota_{1,2}^!(\tilde{\mathcal{E}}) \in D_{Z_1 \times Z_2}(\mathbf{Qcoh}(X_1 \times X_2))$, we have

$$H^0(\mathbf{F}_2^{\mathrm{dg}}) \cong \iota_2^! \circ \iota_2 \circ H^0(\mathbf{F}_2^{\mathrm{dg}}) \circ \iota_1^! \circ \iota_1 \cong \iota_2^! \circ \Phi_{\tilde{\mathcal{E}}} \circ \iota_1 \cong \Phi_{\mathcal{E}}^s$$

by Lemma 2.6.

Remark 4.13. The careful reader may notice that Toën's result [32, Thm. 8.9] was originally proved by using different enhancements for $D(\mathbf{Qcoh}(X_1))$ and $D(\mathbf{Qcoh}(X_2))$. That the same statement holds true for $D^{\mathrm{dg}}(X_1)$ and $D^{\mathrm{dg}}(X_2)$ is observed in [22, Thm. 4.9].

Putting this together with the isomorphisms (4.6) and (4.7), we have proved Proposition 4.10. By Example 2.11, the following consequence of Proposition 4.10 is immediate.

Corollary 4.14. *Let X_1 be a quasi-projective scheme containing a projective subscheme Z_1 such that either X_1 is smooth or $X_1 = Z_1$. Assume that X_2 is a scheme containing a closed subscheme Z_2 . Then, for any exact functor $\mathbf{F}: \mathbf{Perf}_{Z_1}(X_1) \rightarrow \mathbf{Perf}_{Z_2}(X_2)$ satisfying (*) there exist $\mathcal{E} \in D_{Z_1 \times Z_2}(\mathbf{Qcoh}(X_1 \times X_2))$ and an isomorphism of exact functors $\mathbf{F} \cong \Phi_{\mathcal{E}}^s$.*

4.4. The category generated by a spherical object. Let us start with a detour about the derived category of a smooth quasi-projective curve with support condition on a closed point for which we can prove variants of Theorems 1.1 and 1.2. Notice that in this case it is not true that the maximal 0-dimensional torsion subsheaf of \mathcal{O}_Z is trivial. In particular, it is easy to see that the construction in Section 2.2 does not provide an almost ample set. Thus we need a particular treatment that, unfortunately, works only for points embedded in curves.

Proposition 4.15. *Let p be a closed point in a smooth quasi-projective curve C .*

(i) *Let X be a scheme with a closed subscheme Z and let*

$$\mathbf{F}: D_p^b(C) \longrightarrow \mathbf{Perf}_Z(X)$$

be an exact functor such that

$$(4.9) \quad \mathrm{Hom}_{\mathbf{Perf}_Z(X)}(\mathbf{F}(\mathcal{A}), \mathbf{F}(\mathcal{B})[k]) = 0,$$

for all $\mathcal{A}, \mathcal{B} \in \mathbf{Coh}_p(C)$ and all $k < 0$. Then there exist $\mathcal{E} \in D_{\{p\} \times Z}^b(\mathbf{Qcoh}(C \times X))$ and an isomorphism of exact functors $\mathbf{F} \cong \Phi_{\mathcal{E}}^s$.

(ii) *The triangulated category $D_p^b(C)$ has a strongly unique enhancement.*

Proof. By Proposition 2.12 the subcategory \mathbf{C} of $\mathbf{Coh}_p(C)$, whose objects are $\{\mathcal{O}_{np} : n > 0\}$, satisfies property (1) in Definition 2.9. In particular, looking carefully at the construction in Section 4.3, this together with (4.9) is enough to provide an $\mathcal{E} \in D_{\{p\} \times Z}^b(\mathbf{Qcoh}(C \times X))$ and an isomorphism

$$\theta: \mathbf{F}|_{\mathbf{C}} \xrightarrow{\sim} \Phi_{\mathcal{E}}^s|_{\mathbf{C}}.$$

To be precise, the fact that \mathcal{E} is a bounded complex is a consequence of Lemma 5.3 below.

Set \mathbf{D}_0 to be the (strictly) full subcategory of $D_p^b(X)$ whose objects are isomorphic to shifts of objects of $\mathbf{Coh}_p(X)$. By Corollary 3.6, the isomorphism θ extends (uniquely) to an isomorphism compatible with shifts

$$\theta_0: F|_{\mathbf{D}_0} \xrightarrow{\sim} \Phi_{\mathcal{E}}^s|_{\mathbf{D}_0}.$$

Being C a smooth curve, any object $\mathcal{F} \in D_p^b(C)$ can be written (in an essentially unique way) as a finite direct sum of objects of \mathbf{D}_0 . Thus θ_0 extends (uniquely) to the desired isomorphism $F \xrightarrow{\sim} \Phi_{\mathcal{E}}^s$, proving (i).

As for (ii), the same line of reasoning as in the proof of Theorem 1.2 in Section 4.2 works with the only difference that the extension of the isomorphism (4.4) takes place due to Corollary 3.6 instead of Corollary 3.8. Then we argue as in the last part of the proof of (i), i.e. using that any object in $D_p^b(C)$ can be (uniquely) written as a finite direct sum of objects of \mathbf{D}_0 . \square

Obviously, if C is a smooth quasi-projective curve, the Serre functor of $D_p^b(C)$ is the shift by one and so $D_p^b(C)$ is a 1-Calabi–Yau category. Moreover, it is easy to verify that the skyscraper sheaf $\mathcal{O}_p \in \mathbf{Coh}_p(C)$ is a 1-spherical object when p is a \mathbb{k} -rational point. Recall that, as we mentioned in the introduction, an object S in a triangulated category \mathbf{T} is *d-spherical* (where d is a positive integer) if $\mathrm{Hom}_{\mathbf{T}}(S, S[i])$ is trivial if $i \neq 0, d$, while it is isomorphic to \mathbb{k} otherwise.

Remark 4.16. Let \mathbf{T} be an idempotent complete algebraic triangulated category (see [22] for its definition) and let $S \in \mathbf{T}$ be a d -spherical object. In this case, up to equivalence, the full triangulated subcategory \mathbf{T}_S of \mathbf{T} classically generated by S depends neither on the triangulated category \mathbf{T} nor on the d -spherical object S (see [23, Thm. 2.1]).

To prove Proposition 1.3, take S to be a 1-spherical object in an idempotent complete algebraic triangulated category \mathbf{T} . If $C = \mathbb{A}^1$ and p is the origin, we observed before that \mathcal{O}_p is a 1-spherical object and hence, by Remark 4.16, there exists an exact equivalence $\mathbf{T}_S \cong D_p^b(C)$. Now it is enough to apply Proposition 4.15.

5. UNIQUENESS OF FOURIER–MUKAI KERNELS

For functors satisfying (*) and with X_1 and X_2 smooth, the uniqueness of Fourier–Mukai kernels is proved via a direct computation in Section 5.2. As a preliminary step, we study some basic properties of Fourier–Mukai functors in the supported setting. In particular, in Lemma 5.3 we show that the Fourier–Mukai kernels have bounded cohomology. We also make clear that, in general, one cannot expect the kernel to have coherent cohomology.

5.1. Basic properties. Let X_1 and X_2 be schemes containing closed subschemes Z_1 and Z_2 . As explained in the following example, we cannot expect that in general the Fourier–Mukai kernel \mathcal{E} of a functor $\Phi_{\mathcal{E}}: \mathbf{Perf}_{Z_1}(X_1) \rightarrow \mathbf{Perf}_{Z_2}(X_2)$ has (bounded and) coherent cohomology when $Z_i \neq X_i$.

Example 5.1. Suppose that there exists $\mathcal{E} \in D_{Z_1 \times Z_1}^b(X_1 \times X_1)$ such that

$$\Phi_{\mathcal{E}}^s \cong \mathrm{id}: \mathbf{Perf}_{Z_1}(X_1) \rightarrow \mathbf{Perf}_{Z_1}(X_1).$$

By [31, Lemma 7.40], there exist $n > 0$ and $\mathcal{E}_n \in D^b(nZ_1 \times nZ_1)$ such that $\iota_{1,1}(\mathcal{E}) \cong (i_n \times i_n)_*(\mathcal{E}_n)$, where $i_n: nZ_1 \rightarrow X_1$ is the embedding. For any $\mathcal{F}_n \in D^b(nZ_1)$, we have

$$(5.1) \quad (i_n)_*(\mathcal{F}_n) \cong \Phi_{\iota_{1,1}(\mathcal{E})}((i_n)_*(\mathcal{F}_n)) \cong (i_n)_* \circ \Phi_{\mathcal{E}_n}(i_n^* \circ (i_n)_*(\mathcal{F}_n)).$$

Now take $X_1 = \mathbb{P}^k$, $Z_1 = \mathbb{P}^{k-1}$ and $\mathcal{F}_n := \mathcal{O}_{nZ_1}(m)$, for $m \in \mathbb{Z}$. An easy calculation shows that $i_n^* \circ (i_n)_*(\mathcal{F}_n) \cong \mathcal{O}_{nZ_1}(m) \oplus \mathcal{O}_{nZ_1}(m-n)[1]$. Hence to have (5.1) verified, we should have either $\Phi_{\mathcal{E}_n}(\mathcal{O}_{nZ_1}(m)) = 0$ or $\Phi_{\mathcal{E}_n}(\mathcal{O}_{nZ_1}(m-n)) = 0$. But the following isomorphisms should hold at the same time

$$\begin{aligned} \Phi_{\mathcal{E}_n}(\mathcal{O}_{nZ_1}(m)) \oplus \Phi_{\mathcal{E}_n}(\mathcal{O}_{nZ_1}(m-n))[1] &\cong \mathcal{O}_{nZ_1}(m), \\ \Phi_{\mathcal{E}_n}(\mathcal{O}_{nZ_1}(m+n)) \oplus \Phi_{\mathcal{E}_n}(\mathcal{O}_{nZ_1}(m))[1] &\cong \mathcal{O}_{nZ_1}(m+n). \end{aligned}$$

If $\Phi_{\mathcal{E}_n}(\mathcal{O}_{nZ_1}(m-n)) = 0$, then from the second isomorphism we would have that $\mathcal{O}_{nZ_1}(m)[1]$ is a direct summand of $\mathcal{O}_{nZ_1}(m+n)$ which is absurd. Thus $\Phi_{\mathcal{E}_n}(\mathcal{O}_{nZ_1}(m)) = 0$. As this holds for all $m \in \mathbb{Z}$, we get a contradiction.

On the other hand, it is easy to find a kernel of the identity functor. Indeed, denoting by $\Delta: X_1 \rightarrow X_1 \times X_1$ the diagonal embedding, setting $\mathcal{O}_\Delta := \Delta_*(\mathcal{O}_{X_1})$, and defining

$$(5.2) \quad \mathcal{I} := \iota_{1,1}^!(\mathcal{O}_\Delta) \in \mathbf{D}_{Z_1 \times Z_1}^b(\mathbf{Qcoh}(X_1 \times X_1))$$

(notice that \mathcal{I} has bounded cohomologies by Lemma 2.3), we have the following result.

Lemma 5.2. *There exists an isomorphism of exact functors*

$$\mathrm{id} \cong \Phi_{\mathcal{I}}^s: \mathbf{D}_{Z_1}^b(\mathbf{Qcoh}(X_1)) \rightarrow \mathbf{D}_{Z_1}^b(\mathbf{Qcoh}(X_1)).$$

Proof. Indeed, $\Phi_{\mathcal{I}}^s \cong \iota_1^! \circ \Phi_{\mathcal{O}_\Delta} \circ \iota_1$ by Lemma 2.6. This is enough to conclude, since $\Phi_{\mathcal{O}_\Delta} \cong \mathrm{id}$. \square

The following result proves, in particular, that in Theorem 1.1 the kernel \mathcal{E} is actually in $\mathbf{D}_{Z_1 \times Z_2}^b(\mathbf{Qcoh}(X_1 \times X_2))$.

Lemma 5.3. *If $\mathcal{E} \in \mathbf{D}_{Z_1 \times Z_2}(\mathbf{Qcoh}(X_1 \times X_2))$ is such that $\Phi_{\mathcal{E}}^s(\mathbf{Perf}_{Z_1}(X_1)) \subseteq \mathbf{Perf}_{Z_2}(X_2)$, then $\mathcal{E} \in \mathbf{D}_{Z_1 \times Z_2}^b(\mathbf{Qcoh}(X_1 \times X_2))$. Moreover, if $X_1 = Z_1$, then $\mathcal{E} \in \mathbf{D}_{X_1 \times Z_2}^b(X_1 \times X_2)$.*

Proof. By [31, Thm. 6.8], for $i = 1, 2$, the category $\mathbf{D}_{Z_i}(\mathbf{Qcoh}(X_i))$ has a compact generator $G_i \in \mathbf{Perf}_{Z_i}(X_i)$ (see [5] for the case without support conditions). Moreover, by the explicit description of the compact generator in the proof of [31, Thm. 6.8], one sees that $G_1 \boxtimes G_2$ is a compact generator of $\mathbf{D}_{Z_1 \times Z_2}(\mathbf{Qcoh}(X_1 \times X_2))$ (for the non-supported case, see for example [5, Lemma 3.4.1] and [26, Thm. 3.7]).

By [31, Prop. 6.9], the kernel \mathcal{E} has bounded cohomology if and only if there exists an interval $[a, b] \subset \mathbb{R}$ such that $\mathrm{Hom}(G_1 \boxtimes G_2, \mathcal{E}[k]) = 0$, for any $k \notin [a, b]$. But now

$$\mathrm{Hom}(G_1 \boxtimes G_2, \mathcal{E}[k]) \cong \mathrm{Hom}(\iota_1(G_1) \boxtimes \iota_2(G_2), \iota_{1,2}(\mathcal{E})[k]) \cong \mathrm{Hom}(G_2, \Phi_{\mathcal{E}}^s(G_1^\vee)[k])$$

which is non-trivial only for finitely many $k \in \mathbb{Z}$.

Suppose that $Z_1 = X_1$. Then $\mathcal{E} \in \mathbf{D}_{X_1 \times Z_2}^b(X_1 \times X_2)$ if and only if $\tilde{\mathcal{E}} := \iota_{1,2}(\mathcal{E}) \in \mathbf{D}^b(X_1 \times X_2)$. Since $\iota_2 \circ \Phi_{\mathcal{E}}^s \cong \Phi_{\tilde{\mathcal{E}}} \circ \iota_1 \cong \Phi_{\tilde{\mathcal{E}}}$, the functor $\Phi_{\tilde{\mathcal{E}}}$ sends perfect complexes to perfect complexes. Hence we can assume, without loss of generality, that $Z_i = X_i$, for $i = 1, 2$. Then it follows from [28, Cor. 9.13 (4)], where the assumption that the functor is fully faithful is not used (see also [10, Lemma 4.1]), that $\tilde{\mathcal{E}} \in \mathbf{D}^b(X_1 \times X_2)$. \square

We recall that a functor

$$\mathbf{F}: \mathbf{D}_{Z_1}^b(\mathbf{Qcoh}(X_1)) \longrightarrow \mathbf{D}_{Z_2}^b(\mathbf{Qcoh}(X_2))$$

is *bounded* if there is an interval $[a, b] \subset \mathbb{R}$ such that, for any $\mathcal{A} \in \mathbf{Qcoh}_{Z_1}(X_1)$, we have that if $H^i(\mathbf{F}(\mathcal{A})) \neq 0$, then $i \in [a, b]$. We will need the following lemma.

Lemma 5.4. *Assume that the base field \mathbb{k} is perfect. Then every exact functor $\mathbf{F}: \mathbf{D}_{Z_1}^b(X_1) \rightarrow \mathbf{D}_{Z_2}^b(\mathbf{Qcoh}(X_2))$ or $\mathbf{G}: \mathbf{D}_{Z_1}^b(\mathbf{Qcoh}(X_1)) \rightarrow \mathbf{D}_{Z_2}^b(\mathbf{Qcoh}(X_2))$, commuting with arbitrary direct sums, is bounded.*

Proof. To deal with the first part, observe that, by Proposition 2.8, the category $\mathbf{Coh}_{Z_1}(X_1)$ is generated, as an abelian category, by the image of the natural fully faithful functor $i_*: \mathbf{Coh}(Z_1) \hookrightarrow \mathbf{Coh}_{Z_1}(X_1)$. This means that it is enough to show the boundedness of the functor

$$\mathbf{F}' := \mathbf{F} \circ i_*: \mathbf{D}^b(Z_1) \longrightarrow \mathbf{D}_{Z_2}^b(\mathbf{Qcoh}(X_2)).$$

Now this is a straightforward consequence of [31, Thm. 7.39] (here we need \mathbb{k} to be perfect). Indeed, by this result, there exist a positive integer d and a compact object G in $\mathbf{D}(\mathbf{Qcoh}(Z_1))$ such that any object of $\mathbf{D}^b(Z_1)$ is generated by G in at most d steps by taking cones, shifts, direct summands and finite direct sums. Hence \mathbf{F}' is certainly bounded.

The second part, concerning exact functors between the bounded derived categories of quasi-coherent sheaves, is proved using the same argument. Indeed, one applies again Proposition 2.8 and reduces to studying the boundedness of the exact functor

$$\mathbf{G}': \mathbf{D}^b(\mathbf{Qcoh}(Z_1)) \longrightarrow \mathbf{D}_{Z_2}^b(\mathbf{Qcoh}(X_2)).$$

By [31, Thm. 7.39], there exists a positive integer d such that any object of $\mathbf{D}^b(\mathbf{Qcoh}(Z_1))$ is generated by G in at most d steps by taking cones, shifts, direct summands, finite direct sums and arbitrary multiples of the same object (see [31, Sect. 3.1.1]). As \mathbf{G}' commutes with arbitrary direct sums, this is enough to conclude. \square

5.2. The uniqueness of the Fourier–Mukai kernels. Assume that the base field \mathbb{k} is perfect and that X_1 and X_2 are smooth quasi-projective schemes containing projective subschemes Z_1 and Z_2 . Consider $\mathcal{E} \in \mathbf{D}_{Z_1 \times Z_2}^b(\mathbf{Qcoh}(X_1 \times X_2))$ and observe that, obviously, $\mathcal{E} \cong \iota_{1,2}^!(\tilde{\mathcal{E}})$, where $\tilde{\mathcal{E}} := \iota_{1,2}(\mathcal{E}) \in \mathbf{D}^b(\mathbf{Qcoh}(X_1 \times X_2))$. We get a Fourier–Mukai functor

$$(5.3) \quad \Psi_{\mathcal{E}} := \Phi_{\mathcal{O}_{\Delta} \boxtimes \tilde{\mathcal{E}}}: \mathbf{D}(\mathbf{Qcoh}(X_1 \times X_1)) \longrightarrow \mathbf{D}(\mathbf{Qcoh}(X_1 \times X_2)).$$

Let $\tilde{\mathcal{I}} := \iota_{1,1}(\mathcal{I}) \in \mathbf{D}^b(\mathbf{Qcoh}(X_1 \times X_1))$, where \mathcal{I} is the complex defined in (5.2) (such that $\Phi_{\tilde{\mathcal{I}}}^s \cong \text{id}$ by Lemma 5.2). We first prove the following result.

Lemma 5.5. *We have $\Psi_{\mathcal{E}}(\tilde{\mathcal{I}}) \cong \tilde{\mathcal{E}}$ and $\Psi_{\mathcal{E}}(\mathcal{F} \boxtimes \mathcal{G}) \cong \mathcal{F} \boxtimes \Phi_{\tilde{\mathcal{E}}}(\mathcal{G})$ for every $\mathcal{F}, \mathcal{G} \in \mathbf{D}(\mathbf{Qcoh}(X_1))$.*

Proof. A standard computation shows that, for every $\mathcal{D} \in \mathbf{D}(\mathbf{Qcoh}(X_1 \times X_1))$,

$$\Psi_{\mathcal{E}}(\mathcal{D}) \cong \mathcal{D} \star \tilde{\mathcal{E}} := (p_{1,3})_*(p_{1,2}^*(\mathcal{D}) \otimes p_{2,3}^*(\tilde{\mathcal{E}})),$$

where $p_{i,j}$ denote the obvious projections from $X_1 \times X_1 \times X_2$. The second assertion in the statement is then clear. As for the first one, by Lemma 2.4 we have

$$\tilde{\mathcal{I}} = \iota_{1,1} \circ \iota_{1,1}^! \circ \Delta_*(\mathcal{O}_{X_1}) \cong \Delta_* \circ \iota_1 \circ \iota_1^!(\mathcal{O}_{X_1}).$$

Moreover, it is easy to see that $\Delta_*(\mathcal{F}) \star \tilde{\mathcal{E}} \cong p_1^*(\mathcal{F}) \otimes \tilde{\mathcal{E}}$ for every $\mathcal{F} \in D(\mathbf{Qcoh}(X_1))$. It follows that, using the same notation as in the proof of Lemma 2.6,

$$\Psi_{\mathcal{E}}(\tilde{\mathcal{I}}) \cong p_1^* \circ \iota_1 \circ \iota_1^!(\mathcal{O}_{X_1}) \otimes \tilde{\mathcal{E}} \cong \bar{\iota}_1 \circ \bar{\iota}_1^!(\mathcal{O}_{X_1 \times X_2}) \otimes \tilde{\mathcal{E}} \cong \bar{\iota}_1 \circ \bar{\iota}_1^!(\tilde{\mathcal{E}}) \cong \tilde{\mathcal{E}},$$

again by Lemma 2.4. \square

We can use this to prove the following result which is precisely the uniqueness statement in Theorem 1.1.

Proposition 5.6. *Let X_1, X_2, Z_1 and Z_2 be as above and let $F: D_{Z_1}^b(X_1) \rightarrow D_{Z_2}^b(X_2)$ be an exact functor satisfying (*). Assume that there are $\mathcal{E}_1, \mathcal{E}_2 \in D_{Z_1 \times Z_2}^b(\mathbf{Qcoh}(X_1 \times X_2))$ such that*

$$F \cong \Phi_{\mathcal{E}_1}^s \cong \Phi_{\mathcal{E}_2}^s.$$

Then $\mathcal{E}_1 \cong \mathcal{E}_2$.

Proof. Since $\tilde{\mathcal{I}}$ is bounded, there exists a bounded above complex

$$\mathcal{L}^\bullet := \{\dots \rightarrow \mathcal{L}^j \rightarrow \mathcal{L}^{j+1} \rightarrow \dots \rightarrow \mathcal{L}^n \rightarrow 0\} \in D(\mathbf{Qcoh}(X_1 \times X_1))$$

such that $\mathcal{L}^\bullet \cong \tilde{\mathcal{I}}$ and, for any $j \in \mathbb{Z}$, the sheaf \mathcal{L}^j is of the form $P_j \boxtimes M_j$, where P_j and M_j are (possibly infinite) direct sums of sheaves in $\mathbf{Amp}(Z_1, X_1, H_1)$ (use Proposition 2.12).

Using again that $\tilde{\mathcal{I}}$ is a bounded complex, for $m > 0$ sufficiently large, the stupid truncation in position $-m$

$$\mathcal{M}^\bullet := \{0 \rightarrow \mathcal{L}^{-m} \rightarrow \dots \rightarrow \mathcal{L}^n \rightarrow 0\}$$

of \mathcal{L}^\bullet is such that $\mathcal{M}^\bullet \cong \tilde{\mathcal{I}} \oplus \mathcal{K}[m]$, for some $\mathcal{K} \in \mathbf{Qcoh}(X_1 \times X_1)$. Applying term by term the functor $\Psi_{\mathcal{E}_i}$ in (5.3) to \mathcal{M}^\bullet we get a complex of complexes

$$\Psi_{\mathcal{E}_i}(\mathcal{L}^{-m}) \longrightarrow \dots \longrightarrow \Psi_{\mathcal{E}_i}(\mathcal{L}^n).$$

Due to Lemma 5.5, the choice of m sufficiently large, the assumption (*) and Lemma 3.1, this complex has a unique (up to isomorphism) right convolution

$$\mathcal{A}_i := \tilde{\mathcal{E}}_i \oplus \mathcal{K}_i[m],$$

with $\mathcal{K}_i = \Psi_{\mathcal{E}_i}(\mathcal{K}) \in D^b(\mathbf{Qcoh}(X_1 \times X_2))$.

Applying Lemma 3.2 under the hypothesis (*), we get $\mathcal{A}_1 \cong \mathcal{A}_2$. By Lemma 5.4 (here we use that \mathbb{k} is perfect), the functor $\Psi_{\mathcal{E}_i}$ is bounded and so, for m large enough,

$$\mathrm{Hom}(\tilde{\mathcal{E}}_1, \mathcal{K}_2[m]) \cong \mathrm{Hom}(\mathcal{K}_1[m], \tilde{\mathcal{E}}_2) \cong 0.$$

Hence $\iota_{1,2}(\mathcal{E}_1) = \tilde{\mathcal{E}}_1 \cong \tilde{\mathcal{E}}_2 = \iota_{1,2}(\mathcal{E}_2)$, which is equivalent to $\mathcal{E}_1 \cong \mathcal{E}_2$. \square

Remark 5.7. Following a suggestion of D. Orlov, we can show that if X_1 is a projective scheme such that $T_0(\mathcal{O}_{X_1}) = 0$, X_2 is a scheme and $\Phi_{\mathcal{E}}: \mathbf{Perf}(X_1) \rightarrow D^b(X_2)$ is an exact fully faithful functor, then $\mathcal{E} \in D(\mathbf{Qcoh}(X_1 \times X_2))$ is uniquely determined (up to isomorphism).

Indeed, suppose that there exist $\mathcal{F} \in D(\mathbf{Qcoh}(X_1 \times X_2))$ and an isomorphism $\Phi_{\mathcal{F}} \cong \Phi_{\mathcal{E}}$. Consider the quasi-functors

$$\Phi_{\mathcal{E}}^{\mathrm{dg}}, \Phi_{\mathcal{F}}^{\mathrm{dg}}: \mathbf{Perf}^{\mathrm{dg}}(X_1) \rightarrow D^{\mathrm{dg}}(X_2)$$

corresponding to \mathcal{E} and \mathcal{F} under the bijection given by [32, Thm. 8.9]. Let $\mathbf{C} \subseteq \mathbf{D}^{\mathrm{dg}}(\mathbf{Qcoh}(X_2))$ be the full dg subcategory whose objects are the same as those in the essential image of $\Phi_{\mathcal{E}}$. If $\Psi^{\mathrm{dg}}: \mathbf{C} \rightarrow \mathrm{Perf}^{\mathrm{dg}}(X_1)$ is a quasi-functor which is the inverse of $\Phi_{\mathcal{F}}^{\mathrm{dg}}$ in \mathbf{Hqe} , consider the composition

$$\mathbf{F}^{\mathrm{dg}} := \Psi^{\mathrm{dg}} \circ \Phi_{\mathcal{E}}^{\mathrm{dg}}: \mathrm{Perf}^{\mathrm{dg}}(X_1) \longrightarrow \mathrm{Perf}^{\mathrm{dg}}(X_1)$$

which has the property $\Phi_{\mathcal{O}_{\Delta}} \cong \mathrm{id} \cong H^0(\mathbf{F}^{\mathrm{dg}})$. As in the proof of [28, Cor. 9.13], the dg quasi-functor \mathbf{F}^{dg} extends to $\mathbf{G}^{\mathrm{dg}}: \mathbf{D}^{\mathrm{dg}}(X_1) \longrightarrow \mathbf{D}^{\mathrm{dg}}(X_1)$. On the other hand, by [32, Thm. 8.9], there exists (a unique) $\mathcal{G} \in \mathbf{D}(\mathbf{Qcoh}(X_1 \times X_1))$ such that $\mathbf{G}^{\mathrm{dg}} \cong \Phi_{\mathcal{G}}^{\mathrm{dg}}$. Hence

$$\Phi_{\mathcal{O}_{\Delta}} \cong H^0(\mathbf{F}^{\mathrm{dg}}) \cong \Phi_{\mathcal{G}}$$

and, using for example [10, Thm. 1.2], we get $\mathcal{G} \cong \mathcal{O}_{\Delta}$. Therefore $\mathbf{G}^{\mathrm{dg}} \cong \mathrm{id}$ and so $\Phi_{\mathcal{E}}^{\mathrm{dg}} \cong \Phi_{\mathcal{F}}^{\mathrm{dg}}$. Applying again [32, Thm. 8.9], we deduce $\mathcal{E} \cong \mathcal{F}$.

Notice that the proof above does not work if the functor $\Phi_{\mathcal{E}}$ satisfies $(*)$ in the introduction but it is not fully faithful. Nevertheless we expect the result to be true in this case as well.

Acknowledgements. The authors are most grateful to the referee(s) for the insightful comments which allowed them to improve the quality of the paper substantially. We would like to thank Arend Bayer, Daniel Huybrechts, Emanuele Macrì, Dmitri Orlov and Pawel Sosna for comments on an early version of this paper. Part of this paper was written while P.S. visited the Department of Mathematics of the University of Utah and the Institut Henri Poincaré in Paris whose warm hospitality is gratefully acknowledged. It is a pleasure to thank these institutions and the Istituto Nazionale di Alta Matematica for the financial support.

REFERENCES

- [1] M. R. Ballard, *Equivalences of derived categories of sheaves on quasi-projective schemes*, arXiv:0905.3148.
- [2] M. R. Ballard, *Sheaves on local Calabi–Yau varieties*, arXiv:0801.3499.
- [3] A. Bayer, E. Macrì, *The space of stability conditions on the local projective plane*, *Duke Math. J.* **160** (2011), 263–322.
- [4] A. Bondal, M. Larsen, V. Lunts, *Grothendieck ring of pretriangulated categories*, *Int. Math. Res. Not.* **29** (2004), 1461–1495.
- [5] A. Bondal, M. Van den Bergh, *Generators and representability of functors in commutative and noncommutative geometry*, *Moscow Math. J.* **3** (2003), 1–36.
- [6] T. Bridgeland, *Stability conditions on triangulated categories*, *Ann. Math.* **166** (2007), 317–346.
- [7] A. Canonaco, D. Orlov, P. Stellari, *Does full imply faithful?*, *J. Noncommut. Geom.* **7** (2013), 357–371.
- [8] A. Canonaco, P. Stellari, *Uniqueness of dg enhancements for categories of compact objects*, in preparation.
- [9] A. Canonaco, P. Stellari, *Fourier–Mukai functors: a survey*, EMS Ser. Congr. Rep., Eur. Math. Soc. (2013), 27–60.
- [10] A. Canonaco, P. Stellari, *Non-uniqueness of Fourier–Mukai kernels*, *Math. Z.* **272** (2012), 577–588.
- [11] A. Canonaco, P. Stellari, *Twisted Fourier–Mukai functors*, *Adv. Math.* **212** (2007), 484–503.
- [12] V. Drinfeld, *DG quotients of DG categories*, *J. Algebra*, **272** (2004), 643–691.
- [13] C. Fu, D. Yang, *The Ringel–Hall Lie algebra of a spherical object*, *J. London Math. Soc.* **85** (2012), 511–533.
- [14] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics Vol. 52, Springer-Verlag (1977).
- [15] D. Huybrechts, M. Lehn, *The geometry of moduli spaces of sheaves*, Cambridge Mathematical Library, Cambridge University Press (2010).
- [16] A. Ishii, K. Ueda, H. Uehara, *Stability conditions on A_n -singularities*, *J. Differential Geom.* **84** (2010), 87–126.

- [17] A. Ishii, H. Uehara, *Autoequivalences of derived categories on the minimal resolutions of A_n -singularities on surfaces*, J. Differential Geom. **71** (2005), 385–435.
- [18] M. Kashiwara, P. Schapira, *Categories and sheaves*, Grundlehren der Mathematischen Wissenschaften **332**, Springer-Verlag, Berlin (2006).
- [19] Y. Kawamata, *Equivalences of derived categories of sheaves on smooth stacks*, Am. J. Math. **126** (2004), 1057–1083.
- [20] B. Keller, *Chain complexes and stable categories*, Manus. Math. **67** (1990), 379–417.
- [21] B. Keller, *Derived categories and their uses*, in “Handbook of algebra” Vol. 1, North-Holland, Amsterdam (1996), 671–701.
- [22] B. Keller, *On differential graded categories*, International Congress of Mathematicians Vol. II, Eur. Math. Soc., Zürich (2006), 151–190.
- [23] B. Keller, D. Yang, G. Zhou, *The Hall algebra of a spherical object*, J. London Math. Soc. **80** (2009), 771–784.
- [24] M. Kontsevich, *Homological algebra of Mirror Symmetry*, in: Proceedings of the International Congress of Mathematicians (Zurich, 1994, ed. S.D. Chatterji), Birkhäuser, Basel (1995), 120–139.
- [25] A. Kuznetsov, V. Lunts, *Categorical resolutions of irrational singularities*, arXiv:1212.6170.
- [26] K. Lin, D. Pomerleano, *Global matrix factorizations*, arXiv:1101.5847.
- [27] J. Lipman, *Lectures on local cohomology and duality*, in “Local Cohomology and its Applications” Lect. Notes in Pure and Applied Math. **226**, Marcel Dekker, New York (2001), 39–89.
- [28] V. Lunts, D. Orlov, *Uniqueness of enhancements for triangulated categories*, J. Amer. Math. Soc. **23** (2010), 853–908.
- [29] A. Neeman, *The Grothendieck duality theorem via Bousfield’s techniques and Brown representability*, J. Amer. Math. Soc., **9** (1996), 205–236.
- [30] D. Orlov, *Equivalences of derived categories and K3 surfaces*, J. Math. Sci. **84** (1997), 1361–1381.
- [31] R. Rouquier, *Dimensions of triangulated categories*, J. K-theory **1** (2008), 193–258.
- [32] B. Toën, *The homotopy theory of dg-categories and derived Morita theory*, Invent. Math. **167** (2007), 615–667.

A.C.: DIPARTIMENTO DI MATEMATICA “F. CASORATI”, UNIVERSITÀ DEGLI STUDI DI PAVIA, VIA FERRATA 1, 27100 PAVIA, ITALY

E-mail address: `alberto.canonaco@unipv.it`

P.S.: DIPARTIMENTO DI MATEMATICA “F. ENRIQUES”, UNIVERSITÀ DEGLI STUDI DI MILANO, VIA CESARE SARDINI 50, 20133 MILANO, ITALY

E-mail address: `paolo.stellari@unimi.it`

URL: `http://users.unimi.it/stellari`